


KDR monoclonal antibody, clone CIH-11

Catalog # MAB20846 Size 100 uL

Applications



Western Blot (Tissue lysate)

Western Blot (Tissue lysate) analysis of human placenta lysate.

Specification

Product Description Rabbit monoclonal antibody raised against synthetic peptide of human KDR.

Immunogen A synthetic peptide corresponding to human KDR.

Host Rabbit

Reactivity Human

Form Liquid

Purification Affinity purification

Isotype IgG

Recommend Usage
Immunoprecipitation (1:20)
Western Blot (1:500-1000)
The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).

Storage Instruction Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Tissue lysate)

Western Blot (Tissue lysate) analysis of human placenta lysate.

- Immunoprecipitation

Gene Info — KDR

Entrez GeneID [3791](#)

Protein Accession# [P35968](#)

Gene Name KDR

Gene Alias CD309, FLK1, VEGFR, VEGFR2

Gene Description kinase insert domain receptor (a type III receptor tyrosine kinase)

Omim ID [191306 602089](#)

Gene Ontology [Hyperlink](#)

Gene Summary Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas. [provided by RefSeq]

Other Designations soluble VEGFR2|vascular endothelial growth factor receptor 2

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Focal adhesion](#)

- [VEGF signaling pathway](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Angina Pectoris](#)
- [Anoxia](#)
- [Asthma](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cerebral Hemorrhage](#)
- [Colorectal Neoplasms](#)
- [Coronary Disease](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Toxicity](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Femur Head Necrosis](#)
- [Foot Dermatoses](#)
- [Genetic Predisposition to Disease](#)
- [Gilbert Disease](#)

- [Glioma](#)
- [Hot Flashes](#)
- [Hyperbilirubinemia](#)
- [Hypercholesterolemia](#)
- [Hypersensitivity](#)
- [Hypertension](#)
- [Kidney Failure](#)
- [Kidney Neoplasms](#)
- [Leukemia](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Lymphedema](#)
- [Macular Degeneration](#)
- [Metabolic Syndrome X](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Myocardial Infarction](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Osteonecrosis](#)
- [Osteoporosis](#)
- [Psychomotor Performance](#)
- [Recurrence](#)
- [Sarcoidosis](#)

- [Scleroderma](#)
- [Stroke](#)
- [Uterine Cervical Neoplasms](#)