


KRT6A monoclonal antibody, clone KRT6A/2368

Catalog # MAB21273 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human basal cell carcinoma with KRT6A monoclonal antibody, clone KRT6A/2368 (Cat # MAB21273).

Specification

Product Description	Mouse monoclonal antibody raised against full length recombinant human KRT6A.
Immunogen	Recombinant protein corresponding to full length human KRT6A.
Host	Mouse
Theoretical MW (kDa)	56
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG2a, kappa
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1-2 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS (0.05% BSA, 0.05% sodium azide).
Storage Instruction	Store at 4°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human basal cell carcinoma with KRT6A monoclonal antibody, clone KRT6A/2368 (Cat # MAB21273).

Gene Info — KRT6A

Entrez GeneID [3853](#)

Protein Accession# [P02538](#)

Gene Name KRT6A

Gene Alias CK6A, CK6C, CK6D, K6A, K6C, K6D, KRT6C, KRT6D

Gene Description keratin 6A

Omim ID [148041](#) [167200](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. As many as six of this type II cytokeratin (KRT6) have been identified; the multiplicity of the genes is attributed to successive gene duplication events. The genes are expressed with family members KRT16 and/or KRT17 in the filiform papillae of the tongue, the stratified epithelial lining of oral mucosa and esophagus, the outer root sheath of hair follicles, and the glandular epithelia. This KRT6 gene in particular encodes the most abundant isoform. Mutations in these genes have been associated with pachyonychia congenita. The type II cytokeratins are clustered in a region of chromosome 12q12-q13. [provided by RefSeq]

Other Designations

56 cytoskeletal type II keratin|K6D keratin|cytokeratin 6A|cytokeratin 6C|cytokeratin 6D|keratin 6C|keratin, epidermal type II, K6A|keratin, epidermal type II, K6C|keratin, type II cytoskeletal 6D|type II keratin isoform K6c