


KRT8 monoclonal antibody, clone H1 + TS1

Catalog # MAB21278 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung cancer with KRT8 monoclonal antibody, clone H1 + TS1 (Cat # MAB21278).

Specification

Product Description	Mouse monoclonal antibody raised against human KRT8.
Immunogen	Cytoskeleton preparation containing cytokeratin 8 (H1); Keratin preparation from a human carcinoma (TS1).
Host	Mouse
Theoretical MW (kDa)	52.5
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.

Storage Instruction

Store at -20 to -80°C.
Aliquot to avoid repeated freezing and thawing.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung cancer with KRT8 monoclonal antibody, clone H1 + TS1 (Cat # MAB21278).

- Immunofluorescence
- Flow Cytometry

Gene Info — KRT8

Entrez GeneID [3856](#)

Protein Accession# [P05787](#)

Gene Name KRT8

Gene Alias CARD2, CK8, CYK8, K2C8, K8, KO

Gene Description keratin 8

Omim ID [148060](#) [215600](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene is a member of the type II keratin family clustered on the long arm of chromosome 12. Type I and type II keratins heteropolymerize to form intermediate-sized filaments in the cytoplasm of epithelial cells. The product of this gene typically dimerizes with keratin 18 to form an intermediate filament in simple single-layered epithelial cells. This protein plays a role in maintaining cellular structural integrity and also functions in signal transduction and cellular differentiation. Mutations in this gene cause cryptogenic cirrhosis. [provided by RefSeq]

Other Designations cytokeratin 8|keratin, type II cytoskeletal 8

Disease

- [Alzheimer disease](#)
- [Cerebral Amyloid Angiopathy](#)

- [Chronic Disease](#)
- [Disease Progression](#)
- [Drug-Induced Liver Injury](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [Inflammatory Bowel Diseases](#)
- [Liver Cirrhosis](#)
- [Liver Failure](#)
- [Neuroblastoma](#)
- [Pancreatitis](#)