

MME monoclonal antibody, clone CB-CALLA

Catalog # MAB21295 Size 100 ug

Specification

| | |
|----------------------------|---|
| Product Description | Mouse monoclonal antibody raised against human MME. |
| Immunogen | Human CD10. |
| Host | Mouse |
| Reactivity | Human |
| Form | Liquid |
| Purification | Protein A/G purification |
| Isotype | IgG1, kappa |
| Recommend Usage | Flow Cytometry (0.5-1 ug/million cells in 0.1 mL) Immunofluorescence (0.5-1 ug/mL) The optimal working dilution should be determined by the end user. |
| Storage Buffer | In 10 mM PBS (0.05% BSA and 0.05% azide). |
| Storage Instruction | Store at 2 to 8°C. |
| Note | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |

Applications

- Immunofluorescence
- Flow Cytometry

Gene Info — MME

| | |
|--------------------|---|
| Entrez GeneID | 4311 |
| Protein Accession# | P08473 |
| Gene Name | MME |
| Gene Alias | CALLA, CD10, DKFZp686O16152, MGC126681, MGC126707, NEP |
| Gene Description | membrane metallo-endopeptidase |
| Omim ID | 120520 |
| Gene Ontology | Hyperlink |
| Gene Summary | <p>This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a 100-kD type II transmembrane glycoprotein, exists in a single copy of greater than 45 kb. The 5' untranslated region of this gene is alternatively spliced, resulting in four separate mRNA transcripts. The coding region is not affected by alternative splicing. [provided by RefSeq]</p> |
| Other Designations | atriopeptidase common acute lymphocytic leukemia antigen enkephalinase membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase) membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10) membrane metallo-endopeptidase |

Pathway

- [Hematopoietic cell lineage](#)
- [Renin-angiotensin system](#)

Disease

- [Alzheimer disease](#)
- [Anorexia Nervosa](#)
- [Atherosclerosis](#)
- [Brain Injuries](#)
- [Bulimia](#)

- [Calcinosis](#)
- [Cardiovascular Diseases](#)
- [Coronary Artery Disease](#)
- [Diabetes Complications](#)
- [Genetic Predisposition to Disease](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)
- [Osteoporosis](#)