


CXCR2 monoclonal antibody, clone 5F7A11


Catalog # MAB21385 Size 100 ug

Applications




Western Blot (Transfected lysate)

Western Blot analysis of Lane 1: HEK293 and Lane 2: CXCR2-hlgGfC transfected HEK293 cell lysates with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human cervix cancer with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385).



Enzyme-linked Immunoabsorbent Assay

ELISA analysis with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385).



Flow Cytometry

Flow cytometric analysis of HL-60 cells with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385) (Green). Red: Negative Control.

Specification

Product Description	Mouse monoclonal antibody raised against partial recombinant human CXCR2.
Immunogen	Recombinant protein corresponding to human CXCR2.
Host	Mouse
Theoretical MW (kDa)	40.8
Reactivity	Human
Form	Liquid
Isotype	IgG1
Recommend Usage	ELISA (1:10000) Flow Cytometry (1:200-1:400) Immunocytochemistry (1:100) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:200-1:1000) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of Lane 1: HEK293 and Lane 2: CXCR2-hlgFc transfected HEK293 cell lysates with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human cervix cancer with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385).

- Immunocytochemistry

- Enzyme-linked Immunoabsorbent Assay

ELISA analysis with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385).

- Flow Cytometry

Flow cytometric analysis of HL-60 cells with CXCR2 monoclonal antibody, clone 5F7A11 (Cat # MAB21385) (Green). Red: Negative Control.

Gene Info — CXCR2

Entrez GeneID [3579](#)

Protein Accession# [P25025](#)

Gene Name CXCR2

Gene Alias CD182, CDw128b, CMKAR2, IL8R2, IL8RA, IL8RB

Gene Description chemokine (C-X-C motif) receptor 2

Omim ID [146928](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33-q36. [provided by RefSeq]

Other Designations

C-X-C chemokine receptor type 2|CXC-R2|CXCR-2|CXCR2 gene for IL8 receptor type B|GRO/M GSA receptor|IL-8 receptor type 2|IL-8R B|chemokine (CXC) receptor 2|high affinity interleukin-8 receptor B|interleukin 8 receptor B|interleukin 8 receptor type 2|interl

Pathway

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)

Disease

- [Acquired Immunodeficiency Syndrome](#)
- [Adenocarcinoma](#)
- [Alcoholism](#)
- [Asthma](#)
- [Bacteriuria](#)
- [Behcet Syndrome](#)
- [Biliary Tract Neoplasms](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Conduct Disorder](#)
- [Coronary Artery Disease](#)

- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Helicobacter Infections](#)
- [Hematologic Diseases](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Infant](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Leishmaniasis](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Maduromycosis](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)

- [Multiple Sclerosis](#)
- [Neoplasm Invasiveness](#)
- [Obstetric Labor](#)
- [Occupational Diseases](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Periodontal Diseases](#)
- [Periodontitis](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoma](#)
- [Stomach Neoplasms](#)
- [Tuberculosis](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)