


Mouse Anti-Human (lambda chain) secondary antibody, clone JDC-12 (Phycoerythrin)

Catalog # MAB22584 Size 100 ug

Applications



Fluorescence-linked Immunosorbent Assay

FLISA plate was coated with purified human IgG kappa, IgM kappa, IgA kappa, IgG lambda, IgM lambda, and IgA lambda. Immunoglobulins were detected with serially diluted Mouse Anti-Human (lambda chain) secondary antibody, clone JDC-12 (Phycoerythrin).

Specification

Product Description	Mouse monoclonal antibody raised against human lambda light chain.
Immunogen	Human lambda light chain.
Host	Mouse
Reactivity	Human
Specificity	Human/rhesus lambda.
Form	Liquid
Conjugation	Phycoerythrin
Purification	Precipitation method and/or chromatography purification
Isotype	IgG1, kappa
Recommend Usage	<p>ELISA (≤ 1 ug/mL)</p> <p>Flow Cytometry (≤ 0.1 ug/10^6 cells)</p> <p>The optimal working dilution should be determined by the end user.</p>

Storage Buffer	In PBS (0.09% sodium azide).
Storage Instruction	Store in the dark at 4°C. Do not freeze. Avoid prolonged exposure to light. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Fluorescence-linked Immunosorbent Assay

FLISA plate was coated with purified human IgG kappa, IgM kappa, IgA kappa, IgG lambda, IgM lambda, and IgA lambda. Immunoglobulins were detected with serially diluted Mouse Anti-Human (lambda chain) secondary antibody, clone JDC-12 (Phycoerythrin).

Gene Info — IGL@

Entrez GeneID	3535
Protein Accession#	P0CG04 ; P0DOY2 ; P0DOY3 ; A0M8Q6
Gene Name	IGL@
Gene Alias	IGL, MGC88804
Gene Description	immunoglobulin lambda locus
Gene Ontology	Hyperlink

Gene Summary	Immunoglobulins recognize foreign antigens and initiate immune responses such as phagocytosis and the complement system. Each immunoglobulin molecule consists of two identical heavy chains and two identical light chains. There are two classes of light chains, kappa and lambda. This region represents the germline organization of the lambda light chain locus. The locus includes V (variable), J (joining), and C (constant) segments. During B cell development, a recombination event at the DNA level joins a single V segment with a J segment; the C segment is later joined by splicing at the RNA level. Recombination of many different V segments with several J segments provides a wide range of antigen recognition. Additional diversity is attained by junctional diversity, resulting from the random addition of nucleotides by terminal deoxynucleotidyltransferase, and by somatic hypermutation, which occurs during B cell maturation in the spleen and lymph nodes. Several V segments and three C segments are known to be incapable of encoding a protein and are considered pseudogenes. The locus also includes several non-immunoglobulin genes, many of which are pseudogenes or are predicted by automated computational analysis or homology to other species. [provided by RefSeq]
--------------	--

Other Designations	immunoglobulin lambda gene cluster
--------------------	------------------------------------