


# Anti-Rituximab monoclonal antibody, clone 194D7


Catalog # MAB23083      Size 40 ug

## Applications



### Enzyme-linked Immunoabsorbent Assay

Anti-Rituximab monoclonal antibody, clone 194D7 (Cat # MAB23083) binds with Rituximab. Coating antigen: Rituximab, 1 ug/mL. Anti-Rituximab monoclonal antibody, clone 194D7 dilution start from 1000 ng/mL.  $EC_{50}$  = 1.618 ng/mL. While the antibody does not recognize the human IgG Fc fragment (data not shown).



### Sandwich ELISA

The Sandwich ELISA assay is developed by using Anti-Rituximab monoclonal antibody, clone 137C6 (Cat # MAB23082) and Anti-Rituximab monoclonal antibody, clone 194D7 (Cat # MAB23083) as capture and detection antibody, respectively. The sensitivity of detecting Rituximab is about 30 pg/mL.

## Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against Rituximab. Target gene is CD20.
<b>Immunogen</b>	Rituximab.
<b>Host</b>	Rabbit
<b>Specificity</b>	The product is specific for Rituximab. The antibody is recommended as a detection antibody in a pharmacokinetic (PK) bridging assay with capture antibody : Anti-Rituximab monoclonal antibody, clone 137C6 (Cat # <a href="#">MAB23082</a> ).
<b>Form</b>	Lyophilized

Preparation Method	The antibody is produced from a hybridoma resulting from the fusion of partner and B-lymphocytes obtained from a rabbit immunized with Rituximab.
Purification	Protein A purification
Isotype	IgG1
Recommend Usage	ELISA (0.005-1 ug/mL (Direct/Indirect)) Sandwich ELISA The optimal working dilution should be determined by the end user.
Inhibitory IC50	Non-inhibitory
Storage Buffer	Lyophilized from PBS, pH 7.4 (0.02% sodium azide).
Storage Instruction	Store at -20°C. The lyophilized product remains stable up to 1 year at -20°C from date of receipt. After reconstitution with deionized water (or equivalent) to a final concentration of 0.5 mg/mL, it can be stored for 2-3 weeks at 2-8°C or for up to 12 months at -20°C or below. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Enzyme-linked Immunoabsorbent Assay

Anti-Rituximab monoclonal antibody, clone 194D7 (Cat # MAB23083) binds with Rituximab. Coating antigen: Rituximab, 1 ug/mL. Anti-Rituximab monoclonal antibody, clone 194D7 dilution start from 1000 ng/mL.  $EC_{50}$  = 1.618 ng/mL. While the antibody does not recognize the human IgG Fc fragment (data not shown).

- Sandwich ELISA

The Sandwich ELISA assay is developed by using Anti-Rituximab monoclonal antibody, clone 137C6 (Cat # MAB23082) and Anti-Rituximab monoclonal antibody, clone 194D7 (Cat # MAB23083) as capture and detection antibody, respectively. The sensitivity of detecting Rituximab is about 30 pg/mL.

## Gene Info — MS4A1

Entrez GeneID	<a href="#">931</a>
Gene Name	MS4A1
Gene Alias	B1, Bp35, CD20, LEU-16, MGC3969, MS4A2, S7
Gene Description	membrane-spanning 4-domains, subfamily A, member 1
Omim ID	<a href="#">112210</a>

## Gene Ontology

[Hyperlink](#)

## Gene Summary

This gene encodes a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This gene encodes a B-lymphocyte surface molecule which plays a role in the development and differentiation of B-cells into plasma cells. This family member is localized to 11q12, among a cluster of family members. Alternative splicing of this gene results in two transcript variants which encode the same protein. [provided by RefSeq]

## Other Designations

B-lymphocyte cell-surface antigen B1|CD20 antigen|CD20 receptor

## Pathway

- [Hematopoietic cell lineage](#)

## Disease

- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Lymphoma](#)
- [Neoplasm Recurrence](#)
- [Ovarian cancer](#)