


Anti-Nivolumab monoclonal antibody, clone 6G5 (Biotin)

Catalog # MAB23123 Size 40 ug

Applications



Enzyme-linked Immunoabsorbent Assay

Anti-Nivolumab monoclonal antibody, clone 6G5 (Biotin) (Cat # MAB23123) binds with Nivolumab, Coating antigen: Nivolumab, 1 ug/mL. Anti-Nivolumab monoclonal antibody, clone 6G5 (Biotin) dilution start from 1000 ng/mL. EC_{50} = 3.01 ng/mL. While the antibody does not recognize human IgG Fc fragment (data not shown).

Specification

Product Description	Mouse monoclonal antibody raised against Nivolumab. Target gene is PD-1.
Immunogen	Nivolumab.
Host	Mouse
Specificity	The product is specific for Nivolumab. The antibody is recommended as a detection antibody in a pharmacokinetic (PK) bridging assay with capture antibody Anti-Nivolumab monoclonal antibody, clone 8G6G3D8 (Cat # MAB23122).
Form	Lyophilized
Conjugation	Biotin
Preparation Method	This antibody is produced from a hybridoma resulting from the fusion of partner and B-lymphocytes obtained from a mouse immunized with Nivolumab.
Purification	Protein A purification
Isotype	IgG1

Recommend Usage	ELISA (0.01-0.1 ug/mL (Direct/Indirect)) The optimal working dilution should be determined by the end user.
Inhibitory IC50	Non-inhibitory
Storage Buffer	Lyophilized from PBS, pH 7.4 (1% BSA, 0.02% sodium azide).
Storage Instruction	Store at -20°C on dry atmospher. The lyophilized product remains stable up to 1 year at -20°C from date of receipt. After reconstitution with deionized water (or equivalent) to a final concentration of 0.5 mg/mL, it can be stored for 2-3 weeks at 2-8°C or for up to 12 months at -20°C or below. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Enzyme-linked Immunoabsorbent Assay

Anti-Nivolumab monoclonal antibody, clone 6G5 (Biotin) (Cat # MAB23123) binds with Nivolumab, Coating antigen: Nivolumab, 1 ug/mL. Anti-Nivolumab monoclonal antibody, clone 6G5 (Biotin) dilution start from 1000 ng/mL. EC₅₀ = 3.01 ng/mL. While the antibody does not recognize human IgG Fc fragment (data not shown).

Gene Info — PDCD1

Entrez GeneID	5133
Gene Name	PDCD1
Gene Alias	CD279, PD1, SLEB2, hPD-1, hPD-I
Gene Description	programmed cell death 1
Omim ID	152700 600244
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a cell surface membrane protein of the immunoglobulin superfamily. This protein is expressed in pro-B-cells and is thought to play a role in their differentiation. In mice, expression of this gene is induced in the thymus when anti-CD3 antibodies are injected and large numbers of thymocytes undergo apoptosis. Mice deficient for this gene bred on a BALB/c background developed dilated cardiomyopathy and died from congestive heart failure. These studies suggest that this gene product may also be important in T cell function and contribute to the prevention of autoimmune diseases. [provided by RefSeq]
Other Designations	-

Pathway

- [Cell adhesion molecules \(CAMs\)](#)
- [T cell receptor signaling pathway](#)

Disease

- [Addison Disease](#)
- [Alveolitis](#)
- [Arthritis](#)
- [Autoimmune Diseases](#)
- [Carcinoma](#)
- [Choriocarcinoma](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Genetic Predisposition to Disease](#)
- [Graves Disease](#)
- [Hydatidiform Mole](#)
- [Inflammation](#)
- [Kidney Diseases](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Lupus Erythematosus](#)

- [Lupus Nephritis](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Sclerosis](#)
- [Myocardial Infarction](#)
- [Neovascularization](#)
- [Polyendocrinopathies](#)
- [Spondylitis](#)
- [Uterine Cervical Neoplasms](#)
- [Uveomeningoencephalitic Syndrome](#)
- [Vasculitis](#)
- [Wegener Granulomatosis](#)