


MCM2 (phospho S139) monoclonal antibody, clone B12 (FITC)

Catalog # MAB23381 Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of C6 cells with MCM2 (phospho Ser139) monoclonal antibody, clone B12 (FITC)(Cat # MAB23381). Unstained as negative control (blue) or untreated (red) or treated with staurosporine (green).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human MCM2.
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Ser139 of human phospho MCM2
Host	Rabbit
Reactivity	Human
Form	Liquid
Conjugation	FITC
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (5 μ L/million cells) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% NaN_3 , 0.2% BSA)
Storage Instruction	Store at 4°C. Do not freeze.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of C6 cells with MCM2 (phospho Ser139) monoclonal antibody, clone B12 (FITC)(Cat # MAB23381). Unstained as negative control (blue) or untreated (red) or treated with staurosporine (green).

Gene Info — MCM2

Entrez GeneID [4171](#)

Protein Accession# [P49736](#)

Gene Name MCM2

Gene Alias BM28, CCNL1, CDCL1, D3S3194, KIAA0030, MGC10606, MITOTIN, cdc19

Gene Description minichromosome maintenance complex component 2

Omim ID [116945](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are involved in the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre-RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. This protein forms a complex with MCM4, 6, and 7, and has been shown to regulate the helicase activity of the complex. This protein is phosphorylated, and thus regulated by, protein kinases CDC2 and CDC7. [provided by RefSeq]

Other Designations DNA replication licensing factor MCM2|MCM2 minichromosome maintenance deficient 2, mitotin|cell division cycle-like 1|cyclin-like 1|minichromosome maintenance deficient 2 (mitotin)|nuclear protein BM28

Pathway

- [Cell cycle](#)
- [DNA replication](#)

Disease

- [Genetic Predisposition to Disease](#)
- [Ovarian Neoplasms](#)