

CD40 FITC monoclonal antibody, clone hCD40

Catalog # MAB23610

Size 500 ug

Specification

Product Description	Mouse FITC monoclonal antibody raised against synthetic peptide of human CD40.
----------------------------	--

Immunogen	Activated Human B cells
------------------	-------------------------

Host	Mouse
-------------	-------

Form	Lyophilized
-------------	-------------

Purification	Protein-A
---------------------	-----------

Isotype	IgG1
----------------	------

Recommend Usage	For staining (10 μ L/1,000,000 cells) The optimal working dilution should be determined by the end user.
------------------------	---

Storage Buffer	Lyophilized from 1 mg/mL in PBS
-----------------------	---------------------------------

Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
----------------------------	---

Applications

- SDS-PAGE

Gene Info — CD40

Entrez GeneID	958
----------------------	---------------------

Gene Name	CD40
------------------	------

Gene Alias	Bp50, CDW40, MGC9013, TNFRSF5, p50
-------------------	------------------------------------

Gene Description	CD40 molecule, TNF receptor superfamily member 5
Omim ID	109535 606843
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been found to be essential in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]
Other Designations	B cell surface antigen CD40 B cell-associated molecule CD40 antigen CD40 antigen (TNF receptor superfamily member 5) CD40 type II isoform CD40L receptor OTTHUMP00000031699 OTTHUMP00000031700 nerve growth factor receptor-related B-lymphocyte activation mol

Pathway

- [Allograft rejection](#)
- [Asthma](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Cytokine-cytokine receptor interaction](#)
- [Primary immunodeficiency](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)

- [Autoimmune Diseases](#)
- [Brain Ischemia](#)
- [Bronchiolitis](#)
- [Calcinosis](#)
- [Cardiovascular Diseases](#)
- [Colitis](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Diabetic Angiopathies](#)
- [Disease Progression](#)
- [Drug Eruptions](#)
- [Edema](#)
- [Endometriosis](#)
- [Genetic Predisposition to Disease](#)
- [Graves Disease](#)
- [Hashimoto Disease](#)
- [Hematologic Diseases](#)
- [Hepatitis B](#)
- [Hodgkin Disease](#)
- [Hypertension](#)
- [Infant](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)

- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [Myasthenia Gravis](#)
- [Occupational Diseases](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Rupture](#)
- [Stroke](#)
- [Thyroiditis](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Wegener Granulomatosis](#)
- [Werner syndrome](#)