

C-peptide monoclonal antibody, clone C-PEP-01 (Biotin)

Catalog # MAB5025

Size 100 ug

Specification

Product Description	Mouse monoclonal antibody raised against C-peptide.
Immunogen	Human C-peptide conjugated with BSA.
Host	Mouse
Reactivity	Human
Specificity	This antibody reacts specifically with C-peptide, a part of the Proinsulin molecule. Proinsulin consists of the three parts: C-peptide and two long strands of amino acids (alpha and beta chains; later become linked together to form the Insulin molecule). No cross-reactivity with Insulin or other peptide hormones or proteins was observed.
Form	Liquid
Conjugation	Biotin
Isotype	IgG1
Recommend Usage	Immunohistochemistry (25 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (0.09% sodium azide)
Storage Instruction	Store at 4°C. Do not freeze. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Gene Info — INS

Entrez GeneID [3630](#)

Gene Name INS

Gene Alias ILPR, IRDN

Gene Description insulin

Omim ID [125850 176730](#)

Gene Ontology [Hyperlink](#)

Gene Summary After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into two chains (peptide A and peptide B) that are covalently linked via two disulfide bonds. Binding of this mature form of insulin to the insulin receptor (INSR) stimulates glucose uptake. A variety of mutant alleles with changes in the coding region have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. [provided by RefSeq]

Other Designations OTTHUMP00000011161|OTTHUMP00000011162|OTTHUMP00000196038|proinsulin

Publication Reference

- [A monoclonal antibody applicable for determination of C-peptide of human proinsulin by RIA.](#)

Hilgert I, Stolba P, Kristofova H, Stefanova I, Bendlova B, Lebl M, Horejsi V.

Hybridoma 1991 Jun; 10(3):379.

Application: RIA, C-peptide fragments

Pathway

- [Insulin signaling pathway](#)
- [Maturity onset diabetes of the young](#)
- [mTOR signaling pathway](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Regulation of autophagy](#)

- [Type I diabetes mellitus](#)
- [Type II diabetes mellitus](#)

Disease

- [Addison Disease](#)
- [Adenomatous Polyps](#)
- [Alzheimer disease](#)
- [Amenorrhea](#)
- [Anorexia Nervosa](#)
- [Atherosclerosis](#)
- [Autoimmune Diseases](#)
- [Autoimmune polyglandular syndrome](#)
- [Bile Duct Neoplasms](#)
- [Birth Weight](#)
- [Body Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bulimia](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Colorectal Neoplasms](#)
- [Diabetes](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetic Nephropathies](#)

- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Diseases in Twins](#)
- [Drug Toxicity](#)
- [Edema](#)
- [Gallbladder Neoplasms](#)
- [Gallstones](#)
- [Genetic Predisposition to Disease](#)
- [Hirsutism](#)
- [Hyperandrogenism](#)
- [Hypercholesterolemia](#)
- [Hyperinsulinism](#)
- [Hypertension](#)
- [Hypertriglyceridemia](#)
- [Insulin Resistance](#)
- [Kidney Failure](#)
- [Liver Neoplasms](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)
- [Obesity](#)
- [Oligomenorrhea](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Failure](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)

- [Pancreatic Neoplasms](#)
- [Parkinson disease](#)
- [Polycystic Ovary Syndrome](#)
- [Polyendocrinopathies](#)
- [Pre-Eclampsia](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Puberty](#)
- [Stomach Neoplasms](#)
- [Thrombophilia](#)
- [Thyroid Diseases](#)
- [Thyroiditis](#)
- [Tobacco Use Disorder](#)
- [Weight Gain](#)