

Cd8a monoclonal antibody, clone 53-6.7 (APC)

Catalog # MAB5130 Size 100 ug

Specification	
Product Description	Rat monoclonal antibody raised against native Cd8a.
Immunogen	Native purified Cd8a from mouse spleen cells.
Host	Rat
Theoretical MW (kDa)	32-34
Reactivity	Mouse
Specificity	This antibody recognizes mouse CD8a (32-34 KDa; alpha chain of the CD8 antigen).
Form	Liquid
Conjugation	APC
Concentration	0.1 mg/mL
Isotype	lgG2a
Recommend Usage	Flow Cytometry (3 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.2% BSA, 0.09% sodium azide)
Storage Instruction	Store in the dark at 4°C. Do not freeze. Avoid prolonged exposure to light. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Flow Cytometry



Gene Info — Cd8a	
Entrez GeneID	<u>12525</u>
Gene Name	Cd8a
Gene Alias	BB154331, Ly-2, Ly-35, Ly-B, Lyt-2
Gene Description	CD8 antigen, alpha chain
Gene Ontology	<u>Hyperlink</u>
Gene Summary	0
Other Designations	Lyt-2.1 lymphocyte differentiation antigen (AA at 100)

Publication Reference

Development, alteration and real time dynamics of conjunctiva-associated lymphoid tissue.

 $Siebelmann\ S,\ Gehlsen\ U,\ Huttmann\ G,\ Koop\ N,\ Bolke\ T,\ Gebert\ A,\ Stern\ ME,\ Niederkorn\ JY,\ Steven\ P.$

Plos One 2013 Dec; 8(12):e82355.

Application: IHC, Mouse, Lymphoid tissue

 Suppression of alkali burn-induced corneal neovascularization by dendritic cell vaccination targeting VEGF receptor 2.

Mochimaru H, Usui T, Yaguchi T, Nagahama Y, Hasegawa G, Usui Y, Shimmura S, Tsubota K, Amano S, Kawakami Y, Ishida S. Investigative Ophthalmology & Visual Science 2008 May; 49(5):2172.

Application: Func, Mouse, Mouse T cells

• IL-2 in vivo activities and antitumor efficacy enhanced by an anti-IL-2 mAb.

Kamimura D, Sawa Y, Sato M, Agung E, Hirano T, Murakami M.

Journal of Immunology 2006 Jul; 177(1):306.

Application: Flow Cyt, Mouse, Mouse T cells

Directed antigen delivery as a vaccine strategy for an intracellular bacterial pathogen.

Bouwer HG, Alberti-Segui C, Montfort MJ, Berkowitz ND, Higgins DE.

PNAS 2006 Mar; 103(13):5102.



Product Information

 <u>alpha-Galactosylceramide can act as a nasal vaccine adjuvant inducing protective immune responses against</u> <u>viral infection and tumor.</u>

Ko SY, Ko HJ, Chang WS, Park SH, Kweon MN, Kang CY.

Journal of Immunology 2005 Sep; 175(5):3309.

Application: Flow Cyt, Mouse, Mouse T cells

• Beta2 integrins are required for skin homing of primed T cells but not for priming naive T cells.

Grabbe S, Varga G, Beissert S, Steinert M, Pendl G, Seeliger S, Bloch W, Peters T, Schwarz T, Sunderkotter C, Scharffetter-Kochanek K.

The Journal of Clinical Investigation 2002 Jan; 109(2):183.

Application: Flow Cyt, IHC, Human, Mouse, LN cells, Mouse ears, T cells