


# Cd69 monoclonal antibody, clone H1.2F3 (SPRD)

Catalog # MAB5746 Size 100 ug

## Applications



### Flow Cytometry

Fresh BALB/c spleen cells were activated by incubation for 20 hours with PMA + ionomycin, then stained with either hamster IgG-FITC (as an isotype control) or Cd69 monoclonal antibody, clone H1.2F3 (FITC) (Cat # MAB5742). Large, activated cells were then gated and analyzed on a FACScan™ flow cytometer (BDIS, San Jose, CA).

## Specification

|                            |                                                                                                                     |
|----------------------------|---------------------------------------------------------------------------------------------------------------------|
| <b>Product Description</b> | Hamster monoclonal antibody raised against native Cd69.                                                             |
| <b>Immunogen</b>           | Native purified Cd69 from Y245 mouse dendritic epidermal T cell line.                                               |
| <b>Host</b>                | Hamster                                                                                                             |
| <b>Reactivity</b>          | Mouse                                                                                                               |
| <b>Specificity</b>         | Mouse CD69/Very Early Activation Antigen (VEA), Mr 85 KDa (unreduced).                                              |
| <b>Form</b>                | Liquid                                                                                                              |
| <b>Conjugation</b>         | SPRD                                                                                                                |
| <b>Isotype</b>             | IgG, lambda                                                                                                         |
| <b>Recommend Usage</b>     | Flow Cytometry (0.2 ug/10 <sup>6</sup> cells)<br>The optimal working dilution should be determined by the end user. |
| <b>Storage Buffer</b>      | In PBS (0.09% sodium azide)                                                                                         |

**Storage Instruction**

Store in the dark at 4°C. Do not freeze.  
Avoid prolonged exposure to light.  
Aliquot to avoid repeated freezing and thawing.

**Note**

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Immunohistochemistry (Frozen sections)
- Immunoprecipitation
- Flow Cytometry

Fresh BALB/c spleen cells were activated by incubation for 20 hours with PMA + ionomycin, then stained with either hamster IgG-FITC (as an isotype control) or Cd69 monoclonal antibody, clone H1.2F3 (FITC) (Cat # MAB5742). Large, activated cells were then gated and analyzed on a FACScan™ flow cytometer (BDIS, San Jose, CA).

## Gene Info — Cd69

**Entrez GeneID** [12515](#)

**Gene Name** Cd69

**Gene Alias** 5830438K24Rik, A1452015, AIM, VEA

**Gene Description** CD69 antigen

**Gene Ontology** [Hyperlink](#)

**Other Designations** Early activation antigen CD69|Very Early Activation Antigen

## Publication Reference

- [The mouse CD69 gene. Structure, expression, and mapping to the NK gene complex.](#)

Ziegler SF, Levin SD, Johnson L, Copeland NG, Gilbert DJ, Jenkins NA, Baker E, Sutherland GR, Feldhaus AL, Ramsdell F.  
Journal of Immunology 1994 Feb; 152(3):1228.

- [Very early \(VEA\) and very late \(VLA\) activation antigens have distinct functions in T lymphocyte activation.](#)

Yokoyama WM, Maxfield SR, Shevach EM.

Immunological Reviews 1989 Jun; 109:153.

Application: Flow Cyt, Mouse, Mouse T cells

- [Characterization of a cell surface-expressed disulfide-linked dimer involved in murine T cell activation.](#)

Yokoyama WM, Koning F, Kehn PJ, Pereira GM, Stingl G, Coligan JE, Shevach EM.

Journal of Immunology 1988 Jul; 141(2):369.

Application: Flow Cyt, Mouse , Mouse lymphocytes