

Bioactive

PDGFB (Human) Recombinant Protein

Catalog # P3661

Size 10 ug

Specification

Product Description	Human PDGFB (P01127, 82 a.a. - 190 a.a.) partial recombinant protein expressed in <i>Pichia pastoris</i> .
Sequence	SLGSLTIAEPAMIAECKTRTEVFESISRRLLIDRTNANFLVWPPCVEVQRCSGCCNNRNVQCRPTQVQLRPVQVRKIEIVRKPIFKKATVTLEDHLACKCETVAAARPVT
Host	Yeast
Theoretical MW (kDa)	32 (non-reducing con
Form	Lyophilized
Preparation Method	<i>Pichia pastoris</i> expression system
Purification	Ion exchange column and HPLC reverse phase column
Purity	> 90% by SDS-PAGE and HPLC
Endotoxin Level	< 0.1 ng/ug (1 EU/ug)
Activity	The ED ₅₀ was determined by the dose-dependent proliferation of Balb/c 3T3 cells. The expected ED ₅₀ for this effect is 1.0-3.0 ng/mL, corresponding to 1 x 10 ⁶ IU/mg.
Storage Buffer	Lyophilized from 0.02 M acetic acid, pH 6.5
Storage Instruction	Store at -20°C on dry atmosphere for 2 years. After reconstitution with deionized water, store at 4°C for 1 month or store at -20°C for 6 months. Aliquot to avoid repeated freezing and thawing.

Applications

- Functional Study
- SDS-PAGE

Gene Info — PDGFB

Entrez GeneID	5155
Protein Accession#	P01127
Gene Name	PDGFB
Gene Alias	FLJ12858, PDGF2, SIS, SSV, c-sis
Gene Description	platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog)
Omim ID	190040
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the platelet-derived growth factor family. The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a motif of eight cysteines. This gene product can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 7, at sites where this gene and that for COL1A1 are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans resulting from unregulated expression of growth factor. Two alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq]
Other Designations	PDGF, B chain Platelet-derived growth factor, beta polypeptide (oncogene SIS) becaplermin oncogene SIS platelet-derived growth factor 2 platelet-derived growth factor beta platelet-derived growth factor, B chain v-sis platelet-derived growth factor beta p

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Focal adhesion](#)
- [Gap junction](#)
- [Glioma](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)

- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Renal cell carcinoma](#)

Disease

- [Amyotrophic lateral sclerosis](#)
- [Anoxia](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)
- [Head and Neck Neoplasms](#)
- [Neoplasms](#)
- [Obstetric Labor](#)
- [Ovarian Neoplasms](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Recurrence](#)