


Bioactive

Full-Length

# MAPK10 (Human) Recombinant Protein

Catalog # P4712 Size 100 ug

# **Applications**



### Result of activity analysis

Result of activity analysis

Specification	
Product Description	Human MAPK10 (NM_138980, 1 a.a 426 a.a.) full-length recombinant protein expressed in <i>Escher ichia coli</i> .
Host	Escherichia coli
Theoretical MW (kDa)	48.06
Form	Liquid
Preparation Method	Escherichia coli expression system
Purification	Immobilized metal affinity chromatography
Concentration	0.211 ug/uL



## **Product Information**

Activity	118 pmol/ug x min
Quality Control Testing	2 ug/lane SDS-PAGE Stained with Coomassie Blue
Storage Buffer	In 50 mM Hepes, 100 mM NaCl, pH 7.5. (5 mM DTT, 20% glycerol)
Storage Instruction	Store at -80°C.
	Aliquot to avoid repeated freezing and thawing
Note	Result of activity analysis
	Result of activity analysis

## **Applications**

- Functional Study
- SDS-PAGE

Gene Info — MAPK10	
Entrez GenelD	<u>5602</u>
Protein Accession#	NM_138980
Gene Name	MAPK10
Gene Alias	FLJ12099, FLJ33785, JNK3, JNK3A, MGC50974, PRKM10, p493F12, p54bSAPK
Gene Description	mitogen-activated protein kinase 10
Omim ID	<u>602897</u> <u>606369</u>
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This protein is a neuronal-specific form of c-Jun N-terminal kinases (JNKs). Through its phosphorylation and nuclear localization, this kinase plays regulatory roles in the signaling pathways during neuronal a poptosis. Beta-arrestin 2, a receptor-regulated MAP kinase scaffold protein, is found to interact with, and stimulate the phosphorylation of this kinase by MAP kinase kinase 4 (MKK4). Cyclin-dependent kianse 5 can phosphorylate, and inhibit the activity of this kinase, which may be important in preventing neuronal apoptosis. Four alternatively spliced transcript variants encoding distinct isof orms have been reported. [provided by RefSeq



### **Product Information**

**Other Designations** 

JNK3 alpha protein kinase|MAP kinase|OTTHUMP00000161180|OTTHUMP00000161182|OTT HUMP00000161183|c-Jun N-terminal kinase 3|c-Jun kinase 3|stress activated protein kinase JN K3|stress activated protein kinase beta

### **Pathway**

- Adipocytokine signaling pathway
- Colorectal cancer
- Epithelial cell signaling in Helicobacter pylori infection
- ErbB signaling pathway
- Fc epsilon RI signaling pathway
- Focal adhesion
- GnRH signaling pathway
- Insulin signaling pathway
- MAPK signaling pathway
- Neurotrophin signaling pathway
- Pancreatic cancer
- Pathways in cancer
- Toll-like receptor signaling pathway
- Type II diabetes mellitus
- Wnt signaling pathway

### Disease

HIV Infections