


Bioactive

Full-Length

ABL1 (Human) Recombinant Protein

Catalog # P5778 Size 5 ug

Applications



Result of activity analysis

Result of activity analysis

Specification

Product Description	Human ABL1 (NP_005148.2, 2 a.a. - 1130 a.a.) full length recombinant protein with His tag expressed in Baculovirus infected Sf21 cells.
Host	insect
Theoretical MW (kDa)	126
Form	Liquid
Preparation Method	Baculovirus infected insect cell (Sf21) expression system
Purification	Ni-NTA affinity chromatography
Purity	91 % by SDS-PAGE/CBB staining

Activity	The activity was measured by off-chip mobility shift assay. The enzyme was incubated with fluorescence-labeled substrate and Mg (or Mn)/ATP. The phosphorylated and unphosphorylated substrates were separated and detected by LabChip™3000. Substrate : ABLtide. ATP: 100 μM.
Quality Control Testing	SDS-PAGE Stained with Coomassie Blue
Storage Buffer	In 50 mM Tris-HCl, 150 mM NaCl, pH 7.5 (0.05% Brij35, 1 mM DTT, 10% glycerol)
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Result of activity analysis Result of activity analysis

Applications

- Functional Study
- SDS-PAGE

Gene Info — ABL1

Entrez GeneID	25
Protein Accession#	NP_005148.2
Gene Name	ABL1
Gene Alias	ABL, JTK7, bcr/abl, c-ABL, p150, v-abl
Gene Description	c-abl oncogene 1, receptor tyrosine kinase
Omim ID	189980
Gene Ontology	Hyperlink
Gene Summary	The ABL1 protooncogene encodes a cytoplasmic and nuclear protein tyrosine kinase that has been implicated in processes of cell differentiation, cell division, cell adhesion, and stress response. Activity of c-Abl protein is negatively regulated by its SH3 domain, and deletion of the SH3 domain turns ABL1 into an oncogene. The t(9;22) translocation results in the head-to-tail fusion of the BCR (MIM:151410) and ABL1 genes present in many cases of chronic myelogenous leukemia. The DNA-binding activity of the ubiquitously expressed ABL1 tyrosine kinase is regulated by CDC 2-mediated phosphorylation, suggesting a cell cycle function for ABL1. The ABL1 gene is expressed as either a 6- or 7-kb mRNA transcript, with alternatively spliced first exons spliced to the common exons 2-11. [provided by RefSeq]

Other Designations

Abelson murine leukemia viral (v-abl) oncogene homolog 1|OTTHUMP00000022375|OTTHUMP0000022376|bcr/c-abl oncogene protein|proto-oncogene tyrosine-protein kinase ABL1|v-abl Abelson murine leukemia viral oncogene homolog 1

Pathway

- [Axon guidance](#)
- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [ErbB signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Pathogenic Escherichia coli infection - EHEC](#)
- [Pathways in cancer](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Diabetes Complications](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [HIV Infections](#)
- [Leukemia](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)

- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)