

Bioactive

BMP4 (Human) Recombinant Protein

Catalog # P6241

Size 100 ug

Applications

Result of activity analysis

Result of activity analysis

Specification

Product Description	Human BMP4 (P12644) recombinant protein expressed in CHO cells.
Sequence	SPKHHSQRARKKNKNCRRHSLYDFSDVGWNDWVAPPGYQAFYCHGDCPFPLADHLNSTNHAI VQTLVNSVNSSIPKACCVPTELSAISMLYLDEYDKVVLKNYQEMVVEGCGCR
Host	Mammals
Theoretical MW (kDa)	13.1/26.2
Form	Lyophilized
Purity	>= 95%
Endotoxin Level	<= 1 EUs/ug (Kinetic LAL) (50% confidence)
Activity	ED ₅₀ <= 15 ng/mL Alkaline phosphatase activity induced in ATDC-5 cells The values provided above are minimum expected values to pass internal requirements.
Quality Control Testing	Reducing and Non-Reducing SDS PAGE
Conformation	Dimer

Storage Buffer	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA).
Storage Instruction	Stored at -20°C to -80°C for 12 month. After reconstitution with sterile water at 0.1 mg/mL, store at -20°C to -80°C for 3 months, store at 4°C for 1 month. Aliquot to avoid repeated freezing and thawing.
Note	Result of activity analysis Result of activity analysis

Applications

- Western Blot
- Functional Study

Gene Info — BMP4

Entrez GeneID	652
Protein Accession#	P12644
Gene Name	BMP4
Gene Alias	BMP2B, BMP2B1, MCOPS6, ZYME
Gene Description	bone morphogenetic protein 4
Omim ID	112262
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily. The superfamily includes large families of growth and differentiation factors. Bone morphogenetic proteins were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal site. This particular family member plays an important role in the onset of endochondral bone formation in humans, and a reduction in expression has been associated with a variety of bone diseases, including the heritable disorder Fibrodysplasia Ossificans Progressiva. Alternative splicing in the 5' untranslated region of this gene has been described and three variants are described, all encoding an identical protein. [provided by RefSeq]
Other Designations	bone morphogenetic protein 2B

Pathway

- [Basal cell carcinoma](#)
- [Cytokine-cytokine receptor interaction](#)
- [Hedgehog signaling pathway](#)
- [Pathways in cancer](#)
- [TGF-beta signaling pathway](#)

Disease

- [Alzheimer disease](#)
- [Anemia](#)
- [Bone Diseases](#)
- [Calcinosis](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Coloboma](#)
- [Colorectal Neoplasms](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetic Nephropathies](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Gaucher disease](#)
- [Genetic Predisposition to Disease](#)
- [Hemochromatosis](#)

- [Hypospadias](#)
- [Kidney Failure](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Metabolic Syndrome X](#)
- [Microphthalmos](#)
- [Neoplasms](#)
- [Neural Tube Defects](#)
- [Obesity](#)
- [Osteoporosis](#)
- [Ovarian Failure](#)
- [Ovarian Neoplasms](#)
- [Polycystic Ovary Syndrome](#)
- [Prenatal Exposure Delayed Effects](#)
- [Puberty](#)
- [Skin Neoplasms](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)