

Bioactive


HuPro®

MUC-1 (Human) Recombinant Protein

Catalog # P6772

Size 100 ug

Applications



MUC-1 (Human) Recombinant Protein (Cat #P6772) was determined by SDS-PAGE with Coomassie Blue, showing a band at 60-75 kDa.

Result of activity analysis

Result of activity analysis

Specification

Product Description	Human MUC-1 (NP_001018016.1, 1 a.a. - 167 a.a.) partial recombinant protein with Fc and His tag at C-terminus expressed in HEK293 cells.
Host	Human
Theoretical MW (kDa)	60-75
Form	Lyophilized
Preparation Method	Mammalian cell (HEK293) expression system
Purification	Ni-sepharose purification
Purity	> 97% by SDS-PAGE

Endotoxin Level	< 0.1 EU/ug
Activity	Measured by the ability to increase beta-catenin levels in the cytoplasm and nucleus of HCT116 human colon adenocarcinoma cells. 0.01-1 ng/mL of Recombinant Human MUC-1 can effectively increase beta-catenin levels.
Quality Control Testing	SDS-PAGE Stained with Coomassie Blue MUC-1 (Human) Recombinant Protein (Cat #P6772) was determined by SDS-PAGE with Coomassie Blue, showing a band at 60-75 kDa.
Recommend Usage	SDS-PAGE The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from PBS, pH 7.4.
Storage Instruction	Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution to a concentration of 0.1-0.5 mg/mL in sterile distilled water, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Aliquot to avoid repeated freezing and thawing.
Note	Result of activity analysis Result of activity analysis

Applications

- SDS-PAGE

Gene Info — MUC1

Entrez GeneID	4582
Protein Accession#	P15941
Gene Name	MUC1
Gene Alias	CD227, EMA, H23AG, MAM6, PEM, PEMT, PUM
Gene Description	mucin 1, cell surface associated
Omim ID	158340
Gene Ontology	Hyperlink

Gene Summary

This gene is a member of the mucin family and encodes a membrane bound, glycosylated phosphoprotein. The protein is anchored to the apical surface of many epithelia by a transmembrane domain, with the degree of glycosylation varying with cell type. It also includes a 20 aa variable number tandem repeat (VNTR) domain, with the number of repeats varying from 20 to 120 in different individuals. The protein serves a protective function by binding to pathogens and also functions in a cell signaling capacity. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. Multiple alternatively spliced transcript variants that encode different isoforms of this gene have been reported, but the full-length nature of only some has been determined. [provided by RefSeq]

Other Designations

DF3 antigen|H23 antigen|MUC-1/SEC|MUC-1/X|MUC1/ZD|OTTHUMP00000033787|OTTHUMP00000033790|OTTHUMP00000033795|breast carcinoma-associated antigen DF3|episialin|mucin 1|mucin 1, transmembrane|peanut-reactive urinary mucin|polymorphic epithelial mucin|tumor as

Disease

- [Adenocarcinoma](#)
- [Asthma](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Chronic Disease](#)
- [Disease Susceptibility](#)
- [Gastritis](#)
- [Genetic Predisposition to Disease](#)
- [Helicobacter Infections](#)
- [Infant](#)
- [Kidney Failure](#)
- [Lung Diseases](#)
- [Lung Neoplasms](#)

- [Metaplasia](#)
- [NARP](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Stomach Neoplasms](#)