


Bioactive

PDFGB (Human) Recombinant Protein

Catalog # P7140

Size 10 ug

Applications



Result of bioactivity analysis

Result of bioactivity analysis

Specification

Product Description	Human PDGFB (P01127, 82 a.a. - 190 a.a.) partial recombinant protein expressed in <i>Pichia pastoris</i> S.
Sequence	SLGSLTIAEPAMIAECKTRTEVFEISRRLIDRTNANFLVWPPCVEVQRCSGCCNNRNVQCRPTQVQLRPVQVRKIEIVRKKPIFKKATVTLEDHLACKCETVAAARPVTRSPGGGSQEQR
Host	Yeast
Theoretical MW (kDa)	24.3
Form	Lyophilized
Preparation Method	<i>Pichia pastoris</i> expression system
Purity	> 97% by SDS-PAGE
Endotoxin Level	< 1 EU per 1 ug of protein (determined by gel clotting method)
Activity	The ED ₅₀ was determined by the dose-dependent stimulation of proliferation of Balb/c 3T3 cells is 5.0 - 10.0 ng/ml, corresponding to a specific activity of > 3.3 x 10 ⁵ units/mg.
Storage Buffer	Lyophilized from sterile distilled Water up to 100 ug/ml

Storage Instruction

Store at 4°C to 8°C for 1 week. For long term storage store at -20°C to -80°C.
Aliquot to avoid repeated freezing and thawing.

Note

Result of bioactivity analysis
Result of bioactivity analysis

Applications

- Functional Study
- SDS-PAGE

Gene Info — PDGFB

Entrez GeneID

[5155](#)

Protein Accession#

[P01127](#)

Gene Name

PDGFB

Gene Alias

FLJ12858, PDGF2, SIS, SSV, c-sis

Gene Description

platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog)

Omim ID

[190040](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the platelet-derived growth factor family. The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a motif of eight cysteines. This gene product can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 7, at sites where this gene and that for COL1A1 are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans resulting from unregulated expression of growth factor. Two alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq]

Other Designations

PDGF, B chain|Platelet-derived growth factor, beta polypeptide (oncogene SIS)|becaplermin|oncogene SIS|platelet-derived growth factor 2|platelet-derived growth factor beta|platelet-derived growth factor, B chain|v-sis platelet-derived growth factor beta p

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Focal adhesion](#)
- [Gap junction](#)
- [Glioma](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Renal cell carcinoma](#)

Disease

- [Amyotrophic lateral sclerosis](#)
- [Anoxia](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)
- [Head and Neck Neoplasms](#)

- [Neoplasms](#)
- [Obstetric Labor](#)
- [Ovarian Neoplasms](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Recurrence](#)