


TNFSF9 (Human) Recombinant Protein

Catalog # P7660 Size 1 mg

Applications



Specification

Product Description	Human TNFSF9 (P41273, 71 a.a. - 254 a.a.) partial recombinant protein with His tag at C-terminus expressed in <i>Escherichia coli</i> .
Sequence	REGPELSPDDPAGLLDLRQGMFAQLVAQNVLLIDGPLSWYSDPGLAGVSLTGGLSYKEDTKELV VAKAGVYYVFFQLELRRV VAGEGSGSVSLALHLQPLRSAAGAAALALTVDLPPASSEARNSAFG FQGRLLHLSAGQRLGVHLHTEARARHAWQLTQGATVLGLFRVTPEIPAGLPSRSE
Host	<i>Escherichia coli</i>
Theoretical MW (kDa)	16-20
Form	Lyophilized
Preparation Method	<i>Escherichia coli</i> expression system
Purity	> 95% by SDS-PAGE
Endotoxin Level	< 1 EU per 1 ug of protein (determined by LAL method)
Quality Control Testing	SDS-PAGE under reducing condition
Storage Buffer	Lyophilized from sterile distilled Water up to 100 ug/mL
Storage Instruction	Store at 4°C to 8°C for 1 week. For long term storage store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- SDS-PAGE

Gene Info — TNFSF9

Entrez GeneID	8744
Protein Accession#	P41273
Gene Name	TNFSF9
Gene Alias	4-1BB-L, CD137L
Gene Description	tumor necrosis factor (ligand) superfamily, member 9
Omim ID	606182
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This transmembrane cytokine is a bidirectional signal transducer that acts as a ligand for TNFRSF9/4-1BB, which is a costimulatory receptor molecule in T lymphocytes. This cytokine and its receptor are involved in the antigen presentation process and in the generation of cytotoxic T cells. The receptor TNFRSF9/4-1BB is absent from resting T lymphocytes but rapidly expressed upon antigenic stimulation. The ligand encoded by this gene, TNFSF9/4-1BBL, has been shown to reactivate anergic T lymphocytes in addition to promoting T lymphocyte proliferation. This cytokine has also been shown to be required for the optimal CD8 responses in CD8 T cells. This cytokine is expressed in carcinoma cell lines, and is thought to be involved in T cell-tumor cell interaction.
Other Designations	homolog of mouse 4-1BB-L receptor 4-1BB ligand

Pathway

- [Cytokine-cytokine receptor interaction](#)

Disease

- [Birth Weight](#)

- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Leukemia](#)
- [Lymphoproliferative Disorders](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Ovarian Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)