

Inorganic Pyrophosphatase (Yeast)

Catalog # P9371 Size 500 uL

| | | | 4 | |
|--------|-----|------|-----|-----|
| _ | pec | ודוי | cat | ıor |
| \sim | DCC | лп | vai | IUI |
| | | | | |

| Dr | \sim | 100 | CILI | atian |
|----|--------|-----|-------|--------------|
| | uuu | 753 | GI II | ption |
| | | | | |

Inorganic Pyrophosphatase (Yeast) is derived from a recombinant E.Coli strain carrying Saccharom yces cerevisiae ppa gene. The enzyme catalyzes the hydrolysis of inorganic pyrophosphate to form orthophosphate. $P_2O_7^{-4} + H_2O + PPase 2HPO_4^{-2}$ Inorganic Pyrophosphatase (yeast) could hydrolyz e the inorganic pyrophosphate generated by nucleic acid amplification, in vitro transcription and othe reactions, removing the inhibition of the inorganic pyrophosphate generated on the reaction system. The removal of pyrophosphate can shift the reaction equilibrium to the end of product formation, which is beneficial to increase the yield of synthetic product. RNA and DNA synthesis are examples of reactions that can be pulled far in the synthesis direction by the action of inorganic pyrophosphatase.

| Theoretical MW (kDa) | 71 | |
|----------------------|---|--|
| Form | Liquid | |
| Concentration | 0.1 U/µL | |
| Recommend Usage | In Vitro Transcription The optimal working dilution should be determined by the end user. | |
| Storage Buffer | 20 mM Tris-HCl (pH 8.0), 100 mM KCl, 0.1 mM EDTA, 1 mM dithiothreitol, 50% glycerol | |
| Storage Instruction | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. | |

Applications

In vitro Transcription