#### Bioactive

# CCL14 (Human) Recombinant Protein

Catalog # P9475 Size 2 x 10 ug

| Specification        |  |
|----------------------|--|
| Product Description  | Human CCL14 (Q16627, 22 a.a 93 a.a.) partial recombinant protein expressed in <i>Escherichia coli</i> .  |
| Sequence             | TESSSRGPYHPSECCFTYTTYKIPRQRIMDYYETNSQCSKPGIVFITKRGHS_x005F_x005F_x005F<br>_x000Dx005F_x000Dx000D_VCTNPSDKWVQDYIKDMKEN  |
| Host                 | Escherichia coli   |
| Theoretical MW (kDa) | 8.4  |
| Form                 | Lyophilized  |
| Preparation Method   | Escherichia coli expression system   |
| Purity               | > 97.0% by SDS-PAGE  |
| Activity             | The Biological activity is 5 - 20 ng/mL, was determined by the ability to chemoattract Human monocyt es, corresponding to a specific activity of 50000 - 200000 IU/mg. |
| Recommend Usage      | Biological Activity<br>SDS-PAGE<br>The optimal working dilution should be determined by the end user.  |
| Storage Buffer       | Lyophilized from sterile distilled Water is > 100 ug/mL  |
| Storage Instruction  | Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.                            |

## Applications

- Functional Study
- SDS-PAGE



### **Product Information**

### Gene Info — CCL14

| Entrez GenelD      | <u>6358</u>  |
|--------------------|--|
| Protein Accession# | <u>Q16627</u>  |
| Gene Name          | CCL14  |
| Gene Alias         | CC-1, CC-3, CKb1, HCC-1, HCC-3, MCIF, NCC-2, NCC2, SCYA14, SCYL2, SY14   |
| Gene Description   | chemokine (C-C motif) ligand 14  |
| Omim ID            | <u>601392</u>  |
| Gene Ontology      | <u>Hyperlink</u>   |
| Gene Summary       | This gene, CCL14, is one of several CC cytokine genes clustered on 17q11.2. The CC cytokines are secreted proteins characterized by two adjacent cysteines. The cytokine encoded by this gen e induces changes in intracellular calcium concentration and enzyme release in monocytes. Multip le transcript variants encoding different isoforms have been found for this gene. Read-through tran scripts are also expressed that include exons from the upstream cytokine gene CCL15, and are r epresented as GeneID: 348249. [provided by RefSeq |
| Other Designations | OTTHUMP00000176860 chemokine CC-1 chemokine CC-3 small inducible cytokine subfamily A (Cys-Cys), member 14   |
|                    |  |

## Pathway

- Chemokine signaling pathway
- Cytokine-cytokine receptor interaction

### Disease

- Asthma
- Bronchiolitis
- Genetic Predisposition to Disease
- Infant
- Lupus Erythematosus
- <u>Multiple Sclerosis</u>

😵 Abnova

**Product Information** 

- <u>Respiratory Syncytial Virus Infections</u>
- Tobacco Use Disorder