

CXCL9 (Human) Recombinant Protein

Catalog # P9531

Size 20 ug

Specification

Product Description	Human CXCL9 (Q07325, 23 a.a. - 125 a.a.) partial recombinant protein with His tag at N-terminus expressed in <i>Escherichia coli</i> .
Sequence	MGSSHHHHHHSSGLVPRGSHMGSTPVVRKGRCSISTNQGTIHLQSLKDLKQFAPSPSCEKIEIIA TLKNGVQTCLNPDSADVKELIKKEKQVSQKKKQKNGKKHQKKVLRKSRQSRQKKT
Host	<i>Escherichia coli</i>
Theoretical MW (kDa)	14
Form	Liquid
Preparation Method	<i>Escherichia coli</i> expression system
Purity	> 85.0% by SDS-PAGE
Recommend Usage	Biological Activity SDS-PAGE The optimal working dilution should be determined by the end user.
Storage Buffer	In 20mM Tris-HCl pH 8.0 (0.15 M NaCl and 30% glycerol)
Storage Instruction	Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- SDS-PAGE

Gene Info — CXCL9

Entrez GeneID

[4283](#)

Protein Accession#	Q07325
Gene Name	CXCL9
Gene Alias	CMK, Humig, MIG, SCYB9, crg-10
Gene Description	chemokine (C-X-C motif) ligand 9
Omim ID	601704
Gene Ontology	Hyperlink
Gene Summary	The function of this gene has not been specifically defined; however, it is thought to be involved in T cell trafficking. This gene has been localized to 4q21 with INP10, which is also a member of the chemokine family of cytokines. [provided by RefSeq]
Other Designations	monokine induced by gamma interferon small inducible cytokine B9

Pathway

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Asthma](#)
- [Bronchiolitis](#)
- [Colitis](#)
- [Coronary Artery Disease](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Genetic Predisposition to Disease](#)
- [Infant](#)
- [Inflammatory Bowel Diseases](#)
- [Ovarian Neoplasms](#)

- [Respiratory Syncytial Virus Infections](#)
- [Severe Acute Respiratory Syndrome](#)