


# MKI67IP polyclonal antibody

Catalog # PAB0369


Size 200 uL

## Applications



### Western Blot (Cell lysate)

Western blot analysis of MKI67IP in 30 ugs of HeLa cell lysate using MKI67IP polyclonal antibody (Cat # PAB0369) at 1 : 1000.



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis in adipocytes of human breast carcinoma tissue using MKI67IP polyclonal antibody (Cat # PAB0369) at 1 : 500 dilution.

## Specification

<b>Product Description</b>	Rabbit polyclonal antibody raised against synthetic peptide of MKI67IP.
<b>Immunogen</b>	A mixture of synthetic peptides corresponding to amino acids 144-160 and 276-293 of human MKI67IP.
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Form</b>	Liquid
<b>Recommend Usage</b>	The optimal working dilution should be determined by the end user.

<b>Storage Buffer</b>	In PBS (0.2% gelatin, 0.05% sodium azide)
<b>Storage Instruction</b>	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
<b>Note</b>	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Western Blot (Cell lysate)

Western blot analysis of MKI67IP in 30 ugs of HeLa cell lysate using MKI67IP polyclonal antibody (Cat # PAB0369) at 1 : 1000.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis in adipocytes of human breast carcinoma tissue using MKI67IP polyclonal antibody (Cat # PAB0369) at 1 : 500 dilution.

## Gene Info — MKI67IP

<b>Entrez GeneID</b>	<a href="#">84365</a>
<b>Gene Name</b>	MKI67IP
<b>Gene Alias</b>	NIFK, Nopp34
<b>Gene Description</b>	MKI67 (FHA domain) interacting nucleolar phosphoprotein
<b>Gene Ontology</b>	<a href="#">Hyperlink</a>
<b>Gene Summary</b>	This gene encodes a protein that interacts with the forkhead-associated domain of the Ki-67 antigen. The encoded protein may bind RNA and may play a role in mitosis and cell cycle progression. Multiple pseudogenes exist on chromosomes 5, 10, 12, 15, and 19.[provided by RefSeq]
<b>Other Designations</b>	nucleolar phosphoprotein Nopp34 nucleolar protein interacting with the FHA domain of pKi-67