


ABL1 (phospho Y245) polyclonal antibody

Catalog # PAB0397


Size 400 uL

Applications



Western Blot (Tissue lysate)

Western blot analysis of ABL1 (phospho Y245) polyclonal antibody (Cat # PAB0397) in mouse colon tissue lysate (35 ug/lane). ABL (arrow) was detected using the purified polyclonal antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human hepatocellular carcinoma tissue reacted with ABL1 (phospho Y245) polyclonal antibody (Cat # PAB0397) which was peroxidase-conjugated to the secondary antibody followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. HC = hepatocarcinoma.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic phosphopeptide of ABL1.
Immunogen	Synthetic phosphopeptide (conjugated with KLH) corresponding to residues surrounding Y245 of human ABL1.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Protein G purification

Recommend Usage	Western Blot (1:1000) Immunohistochemistry (1:50-100) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Tissue lysate)

Western blot analysis of ABL1 (phospho Y245) polyclonal antibody (Cat # PAB0397) in mouse colon tissue lysate (35 ug/lane). ABL (arrow) was detected using the purified polyclonal antibody.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human hepatocellular carcinoma tissue reacted with ABL1 (phospho Y245) polyclonal antibody (Cat # PAB0397) which was peroxidase-conjugated to the secondary antibody followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. HC = hepatocarcinoma.

Gene Info — ABL1

Entrez GeneID	25
Protein Accession#	NP_005148:P00519
Gene Name	ABL1
Gene Alias	ABL, JTK7, bcr/abl, c-ABL, p150, v-abl
Gene Description	c-abl oncogene 1, receptor tyrosine kinase
Omim ID	189980
Gene Ontology	Hyperlink

Gene Summary

The ABL1 protooncogene encodes a cytoplasmic and nuclear protein tyrosine kinase that has been implicated in processes of cell differentiation, cell division, cell adhesion, and stress response. Activity of c-Abl protein is negatively regulated by its SH3 domain, and deletion of the SH3 domain turns ABL1 into an oncogene. The t(9;22) translocation results in the head-to-tail fusion of the BCR (MIM:151410) and ABL1 genes present in many cases of chronic myelogenous leukemia. The DNA-binding activity of the ubiquitously expressed ABL1 tyrosine kinase is regulated by CDC2-mediated phosphorylation, suggesting a cell cycle function for ABL1. The ABL1 gene is expressed as either a 6- or 7-kb mRNA transcript, with alternatively spliced first exons spliced to the common exons 2-11. [provided by RefSeq]

Other Designations

Abelson murine leukemia viral (v-abl) oncogene homolog 1|OTTHUMP00000022375|OTTHUMP00000022376|bcr/c-abl oncogene protein|proto-oncogene tyrosine-protein kinase ABL1|v-abl Abelson murine leukemia viral oncogene homolog 1

Publication Reference

- [Presenilin Is Necessary for Efficient Proteolysis through the Autophagy-Lysosome System in a \$\gamma\$ -Secretase-Independent Manner.](#)

Neely KM, Green KN, Laferla FM.

J Neurosci 2011 Feb; 31:2781.

Application: WB, Mouse, Mouse embryonic fibroblasts (MEFs)

- [Chronic myeloid leukaemia: an investigation into the role of Bcr-Abl-induced abnormalities in glucose transport regulation.](#)

Barnes K, McIntosh E, Whetton AD, Daley GQ, Bentley J, Baldwin SA.

Oncogene 2005 May; 24(20):3257.

- [Interpretation of submicroscopic deletions of the BCR or ABL gene should not depend on extra signal-FISH: problems in interpretation of submicroscopic deletion of the BCR or ABL gene with extra signal-FISH.](#)

Kim YR, Cho HI, Yoon SS, Park S, Kim BK, Lee YK, Chun H, Kim HC, Lee DS.

Genes, Chromosomes & Cancer 2005 May; 43(1):37.

- [Identification of mcl-1 as a BCR/ABL-dependent target in chronic myeloid leukemia \(CML\): evidence for cooperative antileukemic effects of imatinib and mcl-1 antisense oligonucleotides.](#)

Aichberger KJ, Mayerhofer M, Krauth MT, Skvara H, Florian S, Sonneck K, Akgul C, Derdak S, Pickl WF, Wacheck V, Selzer E, Monia BP, Moriggl R, Valent P, Sillaber C.

Blood 2004 Dec; 105(8):3303.

Pathway

- [Axon guidance](#)
- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [ErbB signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Pathogenic Escherichia coli infection - EHEC](#)
- [Pathways in cancer](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Diabetes Complications](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [HIV Infections](#)
- [Leukemia](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)