

# Rabbit Anti-Bovine IgG F(c) secondary antibody (Rhodamine)

Catalog # PAB10458

Size 1500 ug

## Specification

|                            |  |
|----------------------------|--|
| <b>Product Description</b> | Rabbit anti-bovine IgG recognizes bovine IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Rhodamine. |
| <b>Immunogen</b>           | Bovine IgG F(c) fragment   |
| <b>Host</b>                | Rabbit   |
| <b>Reactivity</b>          | Bovine   |
| <b>Form</b>                | Lyophilized  |
| <b>Conjugation</b>         | Rhodamine (TRITC)  |
| <b>Purification</b>        | This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.                 |
| <b>Recommend Usage</b>     | The optimal working dilution should be determined by the end user.   |
| <b>Storage Buffer</b>      | Lyophilized from 20 mM KH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7.2   |
| <b>Storage Instruction</b> | Store at 4°C on dry atmosphere.<br>After reconstitution with 1 mL of deionized water, store at -20°C or lower.<br>Aliquot to avoid repeated freezing and thawing.                  |

## Applications

- Immunofluorescence
- Flow Cytometry

## Publication Reference

- [Texas Red, a hydrophilic, red-emitting fluorophore for use with fluorescein in dual parameter flow microfluorometric and fluorescence microscopic studies.](#)

J A Titus, R Haugland, S O Sharrow, D M Segal.

Journal of Immunological Methods 1982 Apr; 50(2):193.