

# Goat Anti-Chicken IgG (H&L) secondary antibody (Biotin)

Catalog # PAB10502

Size 2 mg

## Specification

<b>Product Description</b>	Goat anti-chicken IgG recognizes chicken IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Biotin.
<b>Immunogen</b>	Chicken IgG whole molecule
<b>Host</b>	Goat
<b>Reactivity</b>	Chicken
<b>Form</b>	Lyophilized
<b>Conjugation</b>	Biotin
<b>Purification</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
<b>Conjugation Note</b>	10-20 BAC molecules per goat IgG molecule
<b>Recommend Usage</b>	The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	Lyophilized from 20 mM KH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7.2 (0.01% sodium azide)
<b>Storage Instruction</b>	Store at 4°C on dry atmosphere. After reconstitution with deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Note</b>	Minimum cross reactivity to bovine, goat, guinea pig, hamster, horse, human, mouse, rabbit, rat, sheep.

## Applications

- Western Blot
- Immunohistochemistry

- Immunofluorescence
- Enzyme-linked Immunoabsorbent Assay

## Publication Reference

- [Protein biotinylation.](#)

Bayer EA, Wilchek M.

Methods in Enzymology 2003 Dec; 184:138.