

Goat Anti-Dog IgG F(c) secondary antibody (Alkaline Phosphatase)

Catalog # PAB10531

Size 1 mg

Specification

| | |
|----------------------------|---|
| Product Description | Goat anti-dog IgG recognizes dog IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Alkaline Phosphatase. |
| Immunogen | Dog IgG F(c) fragment |
| Host | Goat |
| Reactivity | Dog |
| Form | Liquid |
| Conjugation | Alkaline Phosphatase (Calf Intestine) |
| Purification | This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities. |
| Recommend Usage | The optimal working dilution should be determined by the end user. |
| Storage Buffer | In 50 mM Tris-Cl, 150 mM NaCl, 1 mM MgCl, 0.1 mM ZnCl ₂ , 50% (v/v) glycerol, pH 8.0 (0.01% sodium azide) |
| Storage Instruction | Store at 4°C. Do not Freeze. |
| Note | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |

Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay

- Dot Blot
- Enzyme Immunoassay

Publication Reference

- [Peroxidase labelled antibody and Fab conjugates with enhanced intracellular penetration.](#)

Avrameas S, Ternynck T.

Immunochemistry 1971 Dec; 8(12):1175.