

Rabbit Anti-Dog IgG F(c) secondary antibody (Fluorescein)

Catalog # PAB10545

Size 1500 ug

Specification

Product Description	Rabbit anti-dog IgG recognizes dog IgG F(c) fragment. This secondary antibody was purified using a ntigen affinity chromatography. The antibody is conjugated with Fluorescein.
Immunogen	Dog IgG F(c) fragment
Host	Rabbit
Reactivity	Dog
Form	Lyophilized
Conjugation	FITC
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Conjugation Note	2.5 moles FITC per mole of rabbit IgG
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% sodium azide)
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunofluorescence
- Flow Cytometry

Publication Reference

- [Conjugation of fluorescein isothiocyanate to antibodies. I. Experiments on the conditions of conjugation.](#)

The TH, Feltkamp TE.

Immunology 1970 Jun; 18(6):865.

Application: Conjugation, Fluorescein isothiocyanate (FITC)