

Rabbit Anti-Dog IgG F(c) secondary antibody (Peroxidase)

Catalog # PAB10546

Size 1500 ug

Specification

| | |
|----------------------------|--|
| Product Description | Rabbit anti-dog IgG recognizes dog IgG F(c) fragment. This secondary antibody was purified using a ntigen affinity chromatography. The antibody is conjugated with Peroxidase. |
| Immunogen | Dog IgG F(c) fragment |
| Host | Rabbit |
| Reactivity | Dog |
| Form | Lyophilized |
| Conjugation | HRP |
| Purification | This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities. |
| Recommend Usage | The optimal working dilution should be determined by the end user. |
| Storage Buffer | Lyophilized from 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% gentamicin sulfate) |
| Storage Instruction | Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |

Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay
- Dot Blot

- Enzyme Immunoassay

Publication Reference

- [Immunohistochemistry with enzyme labeled antibodies: a brief review.](#)

Farr AG, Nakane PK.

J Immunol Methods 1981 Jan; 47(2):129.