

Chicken Anti-Goat IgG (H&L) secondary antibody (Biotin)

Catalog # PAB10566

Size 1 mg

Specification

Product Description	Chicken anti-goat IgG recognizes goat IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Biotin.
Immunogen	Goat IgG whole molecule
Host	Chicken
Reactivity	Goat
Form	Liquid
Conjugation	Biotin
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Conjugation Note	10-20 BAC molecules per chicken IgG molecule
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	In 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry
- Immunofluorescence

- Enzyme-linked Immunoabsorbent Assay

Publication Reference

- [Protein biotinylation.](#)

Bayer EA, Wilchek M.

Methods in Enzymology 2003 Dec; 184:138.