

Rabbit Anti-Goat IgG F(c) secondary antibody (Fluorescein)

Catalog # PAB10587

Size 2 mg

Specification

Product Description	Rabbit anti-goat IgG recognizes goat IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Fluorescein.
Immunogen	Goat IgG F(c) fragment
Host	Rabbit
Reactivity	Goat
Form	Lyophilized
Conjugation	FITC
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Conjugation Note	3.1 moles FITC per mole of rabbit IgG
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 10 mM Sodium Phosphate, 150 mM Sodium Chloride, pH 7.2 (10 mg/mL BSA, 0.01% thimerosal)
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	This product contains thimerosal: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunofluorescence
- Flow Cytometry

Publication Reference

- [Immunofluorescence using dichlorotriazinylaminofluorescein \(DTAF\). I. Preparation and fractionation of labelled IgG.](#)

D Blakeslee, M G Baines.

Journal of Immunological Methods 1976 Dec; 13(3-4):305.