

# Rabbit Anti-Goat IgG F(c) secondary antibody (Peroxidase)

Catalog # PAB10588

Size 2 mg

## Specification

<b>Product Description</b>	Rabbit anti-goat IgG recognizes goat IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Peroxidase.
<b>Immunogen</b>	Goat IgG F(c) fragment
<b>Host</b>	Rabbit
<b>Reactivity</b>	Goat
<b>Form</b>	Lyophilized
<b>Conjugation</b>	HRP
<b>Purification</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
<b>Recommend Usage</b>	The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	Lyophilized from 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 (0.01% gentamicin sulfate, 10 mg/mL BSA).
<b>Storage Instruction</b>	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

## Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay

## Publication Reference

- [Immunohistochemistry with enzyme labeled antibodies: a brief review.](#)

Farr AG, Nakane PK.

J Immunol Methods 1981 Jan; 47(2):129.