

Rabbit Anti-Goat IgG F(c) secondary antibody (Biotin)

Catalog # PAB10590 Size 2 mg

Specification	
- Opcomodion	
Product Description	Rabbit anti-goat IgG recognizes goat IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Biotin.
Immunogen	Goat IgG F(c) fragment
Host	Rabbit
Reactivity	Goat
Form	Lyophilized
Conjugation	Biotin
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% sodium azide)
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry
- Immunofluorescence



• Enzyme-linked Immunoabsorbent Assay

Publication Reference

Protein biotinylation.

Bayer EA, Wilchek M.

Methods in Enzymology 2003 Dec; 184:138.