

# Goat Anti-Golden Syrian hamster IgG (H&L) secondary antibody (Biotin)

Catalog # PAB10596

Size 2 mg

## Specification

|                            |  |
|----------------------------|--|
| <b>Product Description</b> | Goat anti-Golden Syrian hamster IgG recognizes Golden Syrian hamster IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Biotin. |
| <b>Immunogen</b>           | Golden Syrian hamster IgG whole molecule   |
| <b>Host</b>                | Goat   |
| <b>Reactivity</b>          | Hamster  |
| <b>Form</b>                | Lyophilized  |
| <b>Conjugation</b>         | Biotin   |
| <b>Purification</b>        | This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.   |
| <b>Recommend Usage</b>     | The optimal working dilution should be determined by the end user.   |
| <b>Storage Buffer</b>      | Lyophilized from 20 mM KH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7.2 (0.01% sodium azide)  |
| <b>Storage Instruction</b> | Store at 4°C on dry atmosphere.<br>After reconstitution with 1 mL of deionized water, store at -20°C or lower.<br>Aliquot to avoid repeated freezing and thawing.  |
| <b>Note</b>                | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |

## Applications

- Western Blot
- Immunohistochemistry

- Immunofluorescence
- Enzyme-linked Immunoabsorbent Assay

## Publication Reference

- [Protein biotinylation.](#)

Bayer EA, Wilchek M.

Methods in Enzymology 2003 Dec; 184:138.