

Goat Anti-Horse IgG F(c) secondary antibody (Biotin)

Catalog # PAB10658

Size 2 mg

Specification

Product Description	Goat anti-horse IgG recognizes horse IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Biotin.
Immunogen	Horse IgG F(c) fragment
Host	Goat
Reactivity	Horse
Form	Lyophilized
Conjugation	Biotin
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 10 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% thimerosal)
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	This product contains thimerosal: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry
- Immunofluorescence

- Enzyme-linked Immunoabsorbent Assay

Publication Reference

- [Protein biotinylation.](#)

Bayer EA, Wilchek M.

Methods in Enzymology 2003 Dec; 184:138.