

Goat Anti-Human IgG F(c) secondary antibody (Peroxidase)

Catalog # PAB10704

Size 2 mg

Specification

Product Description	Goat anti-human IgG recognizes human IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Peroxidase.
Immunogen	Human IgG F(c) fragment
Host	Goat
Reactivity	Human
Form	Lyophilized
Conjugation	HRP
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	In 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% gentamicin sulfate, 10 mg/mL BSA, Immunoglobulin and Protease free)
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay

Publication Reference

- [Immunohistochemistry with enzyme labeled antibodies: a brief review.](#)

Farr AG, Nakane PK.

J Immunol Methods 1981 Jan; 47(2):129.