

Chicken Anti-Mouse IgG (H&L) secondary antibody (Fluorescein)

Catalog # PAB10733 Size 1 mg

Specification	
Product Description	Chicken anti-mouse IgG recognizes mouse IgG whole molecule. This secondary antibody was purifie d using antigen affinity chromatography. The antibody is conjugated with Fluorescein.
Immunogen	Mouse IgG whole molecule
Host	Chicken
Reactivity	Mouse
Form	Lyophilized
Conjugation	FITC
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities.
Conjugation Note	3.5 moles FITC per mole of chicken lgG
Recommend Usage	Immunofluorescence (1:1000-1:5000) Flow Cytometry (1:500-1:2500) The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 20 mM potassium phosphate, 150 mM NaCl, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin and Protease free))
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Immunofluorescence



Flow Cytometry

Publication Reference

• Conjugation of fluorescein isothiocyanate to antibodies. I. Experiments on the conditions of conjugation.

The TH, Feltkamp TE.

Immunology 1970 Jun; 18(6):865.

Application: Conjugation, Fluorescein isothiocyanate (FITC)