

# Goat Anti-Rabbit IgG F(ab')<sub>2</sub> secondary antibody (Peroxidase)

Catalog # PAB10828

Size 2 mg

## Specification

<b>Product Description</b>	Goat anti-rabbit IgG recognizes rabbit IgG F(ab') <sub>2</sub> fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Peroxidase.
<b>Immunogen</b>	Rabbit IgG F(ab') <sub>2</sub> fragment
<b>Host</b>	Goat
<b>Reactivity</b>	Rabbit
<b>Form</b>	Lyophilized
<b>Conjugation</b>	HRP
<b>Purification</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
<b>Conjugation Note</b>	
<b>Recommend Usage</b>	ELISA (1:10000-1:50000) Immunohistochemistry (1:500-1:2500) Western Blot (1:1000-1:10000) The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	Lyophilized from 0.01 M sodium phosphate, 0.15 M sodium chloride, pH 7.2 (0.01% (w/v) gentamicin sulfate (Do NOT add sodium azide!)), 10 mg/mL BSA (Immunoglobulin and Protease free)). Restore with 1 mL deionized water (or equivalent).
<b>Storage Instruction</b>	Store at 4°C prior to restoration. For extended storage aliquot contents and freeze at -20°C or below. Aliquot to avoid repeated freezing and thawing.

## Applications

- Western Blot
- Immunohistochemistry

- Enzyme-linked Immunoabsorbent Assay
- Dot Blot
- Enzyme Immunoassay

## Publication Reference

- [Immunohistochemistry with enzyme labeled antibodies: a brief review.](#)

Farr AG, Nakane PK.

J Immunol Methods 1981 Jan; 47(2):129.