

Goat Anti-Rat IgG F(c) secondary antibody (Peroxidase)

Catalog # PAB10864

Size 2 mg

Specification

Product Description Goat anti-rat IgG recognizes rat IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Peroxidase.

Immunogen Rat IgG F(c) fragment

Host Goat

Reactivity Rat

Form Lyophilized

Conjugation HRP

Purification This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.

Conjugation Note

Recommend Usage The optimal working dilution should be determined by the end user.

Storage Buffer Lyophilized from 10 mM potassium phosphate, 75 mM NaCl, 50% (v/v) glycerol, pH 7.2 (0.01% (w/v) gentamicin sulfate (Do NOT add sodium azide!), 10 mg/mL BSA (Immunoglobulin and Protease free))

Storage Instruction Store at 4°C on dry atmosphere.
After reconstitution with 1 mL of deionized water, store at -20°C or lower.
Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay

- Dot Blot
- Enzyme Immunoassay

Publication Reference

- [Immunohistochemistry with enzyme labeled antibodies: a brief review.](#)

Farr AG, Nakane PK.

J Immunol Methods 1981 Jan; 47(2):129.