

Goat Anti-Rat IgG F(ab')2 secondary antibody (Peroxidase)

Catalog # PAB11077 Size 500 uL

Specification	
Product Description	Goat anti-rat IgG recognizes rat IgG F(ab')2 fragment. This secondary antibody was purified using an tigen affinity chromatography. The antibody is conjugated with Peroxidase.
Immunogen	Rat lgG F(ab')2 fragment
Host	Goat
Reactivity	Rat
Specificity	No reaction was observed against anti-Pepsin, anti-goat IgG F(c), rat IgG F(c) or bovine, horse and h uman serum proteins.
Form	Lyophilized
Conjugation	HRP
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followe d by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatog raphic separation.
Conjugation Note	
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 10 mM Sodium Phosphate, 25 mM Sodium Chloride, pH 7.2 (15 mg/mL BSA, Imm unoglobulin and Protease free)
Storage Instruction	Store at -20°C. After reconstitution with 500 uL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	Minimum cross reactivity to bovine, horse, human.

Applications

Western Blot



- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay