


# TAC3 polyclonal antibody

Catalog # PAB12010 Size 100 uL

## **Applications**



## Western Blot (Tissue lysate)

Western blot analysis of TAC3 in rat brain tissue with TAC3 polyclonal antibody (Cat # PAB12010).

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic peptide of TAC3.
Immunogen	A synthetic peptide corresponding to mouse TAC3.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Form	Liquid
Quality Control Testing	Antibody Reactive Against Synthetic Peptide.
Recommend Usage	Immunohistochemistry (1:500-1:1000)  The optimal working dilution should be determined by the end user.
Storage Buffer	In Tris-citrate/phosphate buffer, pH 7.0-8.0 (0.01% sodium azide)
Storage Instruction	Store at -20°C or -80°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.



### **Applications**

Western Blot (Tissue lysate)

Western blot analysis of TAC3 in rat brain tissue with TAC3 polyclonal antibody (Cat # PAB12010).

Immunohistochemistry

### **Publication Reference**

A tachykinin-like factor increases glutamate toxicity in rat cerebellar granule cells.

Severini C, Ciotti MT, Mercanti D, Barbato C, Calissano P.

Neuropharmacology 2003 Jan; 44(1):117.

Application: Incubated, Recombinant protein

 Identification of neurokinin B-expressing neurons as an highly estrogen-receptive, sexually dimorphic cell group in the ovine arcuate nucleus.

Goubillon ML, Forsdike RA, Robinson JE, Ciofi P, Caraty A, Herbison AE.

Endocrinology 2000 Nov; 141(11):4218.

Application: IHC, Sheep, Brain

Neurokinin-3 receptor distribution in rat and human brain: an immunohistochemical study.

Mileusnic D, Lee JM, Magnuson DJ, Hejna MJ, Krause JE, Lorens JB, Lorens SA.

Neuroscience 1999 Jan; 89(4):1269.

Application: C-ELISA, IHC, WB-Ti, WB-Tr, Human, Mouse, Rat, Cortex, CHO cells, Hypothalamus, Recombinant protein