


PCNA polyclonal antibody

Catalog # PAB12722 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil tissue stained with PCNA polyclonal antibody (Cat # PAB12722) at 1 : 200 for 10 min at RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

Specification

Product Description	Rabbit polyclonal antibody raised against recombinant PCNA.
Immunogen	Recombinant protein corresponding to human PCNA.
Host	Rabbit
Reactivity	Human
Specificity	This antibody recognizes human PCNA.
Form	Liquid
Quality Control Testing	Antibody Reactive Against Recombinant Protein.
Recommend Usage	Western Blot (0.1-1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (2-5 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In TBS, pH 7.2 (BSA, 10% Proclin300)
Storage Instruction	Store at -20°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil tissue stained with PCNA polyclonal antibody (Cat # PAB12722) at 1 : 200 for 10 min at RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

Gene Info — PCNA

Entrez GeneID	5111
Gene Name	PCNA
Gene Alias	MGC8367
Gene Description	proliferating cell nuclear antigen
Omim ID	176740
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome. [provided by RefSeq]
Other Designations	DNA polymerase delta auxiliary protein OTTHUMP00000030189 OTTHUMP00000030190 cyclin

Pathway

- [Base excision repair](#)
- [Cell cycle](#)
- [DNA replication](#)
- [Mismatch repair](#)
- [Nucleotide excision repair](#)

Disease

- [Adenocarcinoma](#)
- [Ataxia telangiectasia](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Colorectal Neoplasms](#)
- [Disease Progression](#)
- [DNA Damage](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)
- [Graft vs Host Disease](#)
- [Head and Neck Neoplasms](#)
- [Kidney Failure](#)
- [Laryngeal Neoplasms](#)
- [Lung Neoplasms](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Neuroma](#)
- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Pharyngeal Neoplasms](#)

- [Pulmonary Disease](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Werner syndrome](#)