

# DLX5 polyclonal antibody

Catalog # PAB13670 Size 100 ug

## Applications



### Western Blot (Tissue lysate)

DLX5 polyclonal antibody (Cat # PAB13670) (1 ug/mL) staining of human bone marrow lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

| Specification        |  |
|----------------------|--|
| Product Description  | Goat polyclonal antibody raised against synthetic peptide of DLX5.   |
| Immunogen            | A synthetic peptide corresponding to amino acids at internal region of human DLX5.   |
| Sequence             | C-AYNRVPSATNQPEK   |
| Host                 | Goat   |
| Theoretical MW (kDa) | 31.5   |
| Reactivity           | Human  |
| Specificity          | Approximately 30 KDa band observed in human bone marrow lysates (calculated MW of 31.5 KDa a ccording to NP_005212.1). The observed molecular weight corresponds to earlier findings in literatur e with different antibodies (Lee et al, J Biol Chem. 2003 Sep 5;278(36):34387-94; PMID: 12815054). |
| Form                 | Liquid   |
| Purification         | Antigen affinity purification  |
| Concentration        | 0.5 mg/mL  |



## **Product Information**

| Recommend Usage     | ELISA (1:32000)<br>Western Blot (1-3 ug/mL)<br>The optimal working dilution should be determined by the end user.       |
|---------------------|---|
| Storage Buffer      | In Tris saline, pH 7.3 (0.5% BSA, 0.02% sodium azide)   |
| Storage Instruction | Store at -20°C.<br>Aliquot to avoid repeated freezing and thawing.  |
| Note                | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only. |

# Applications

#### • Western Blot (Tissue lysate)

DLX5 polyclonal antibody (Cat # PAB13670) (1 ug/mL) staining of human bone marrow lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

• Enzyme-linked Immunoabsorbent Assay

| Gene Info — DLX5   |  |
|--------------------|--|
| Entrez GenelD      | 1749   |
| Protein Accession# | <u>NP_005212.1</u>   |
| Gene Name          | DLX5   |
| Gene Alias         | -  |
| Gene Description   | distal-less homeobox 5   |
| Omim ID            | <u>600028</u>  |
| Gene Ontology      | Hyperlink  |
| Gene Summary       | This gene encodes a member of a homeobox transcription factor gene family similiar to the Dros ophila distal-less gene. The encoded protein may play a role in bone development and fracture he aling. Mutation in this gene, which is located in a tail-to-tail configuration with another member of t he family on the long arm of chromosome 7, may be associated with split-hand/split-foot malforma tion. [provided by RefSeq |
| Other Designations | distal-less homeo box 5  |



### **Product Information**

### **Publication Reference**

 DLX5 and HOXC8 enhance the chondrogenic differentiation potential of stem cells from apical papilla via LINC01013.

Haoqing Yang, Yangyang Cao, Jianpeng Zhang, Yuncun Liang, Xiaomin Su, Chen Zhang, Huina Liu, Xiao Han, Lihua Ge, Zhipeng Fan.

Stem Cell Research & Therapy 2020 Jul; 11(1):271.

Application: IP, WB-Tr, Human, Stem cells from apical papilla

 <u>Distal-less homeobox 5 promotes the osteo-/dentinogenic differentiation potential of stem cells from apical</u> papilla by activating histone demethylase KDM4B through a positive feedback mechanism.

Yang H, Fan J, Cao Y, Gao R, Fan Z.

Experimental Cell Research 2019 Jan; 374(1):221.

Application: WB-Ce, WB-Tr, Human, HEK 293T cells, Stem cells from apical papilla

 A novel recurrent chromosomal inversion implicates the homeobox gene DIx5 in T-cell lymphomas from Lck-Akt2 transgenic mice.

Tan Y, Timakhov RA, Rao M, Altomare DA, Xu J, Liu Z, Gao Q, Jhanwar SC, Di Cristofano A, Wiest DL, Knepper JE, Testa JR. Cancer Research 2008 Mar; 68(5):1296.

#### Disease

- Cleft Lip
- <u>Cleft Palate</u>