


SIRPA polyclonal antibody

Catalog # PAB14459 Size 100 ug

Applications



Western Blot (Cell lysate)

Western blot analysis of SIRP alpha in THP-1 cell lysate with SIRPA polyclonal antibody (Cat # PAB14459) at 1:2000 dilution. Bands of molecular weight 75-110 kDa correspond to SIRP alpha 1, 2 and 3.

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic peptide of SIRPA.
Immunogen	A synthetic peptide corresponding to C-terminus of human SIRPA.
Host	Rabbit
Reactivity	Bovine, Human, Mouse, Rat
Form	Liquid
Recommend Usage	Western Blot (1:1000-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.



Applications

Western Blot (Cell lysate)

Western blot analysis of SIRP alpha in THP-1 cell lysate with SIRPA polyclonal antibody (Cat # PAB14459) at 1:2000 dilution. Bands of molecular weight 75-110 kDa correspond to SIRP alpha 1, 2 and 3.

Gene Info — SIRPA	
Entrez GenelD	<u>140885</u>
Gene Name	SIRPA
Gene Alias	BIT, CD172A, MFR, MYD-1, P84, PTPNS1, SHPS-1, SHPS1, SIRP, SIRP-ALPHA-1, SIRPalpha, SIRPalpha2
Gene Description	signal-regulatory protein alpha
Omim ID	<u>602461</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transme mbrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein can be phosphorylated by tyrosine kinases. The phosp ho-tyrosine residues of this PTP have been shown to recruit SH2 domain containing tyrosine phos phatases (PTP), and serve as substrates of PTPs. This protein was found to participate in signal t ransduction mediated by various growth factor receptors. CD47 has been demonstrated to be a li gand for this receptor protein. This gene and its product share very high similarity with several oth er members of the SIRP family. These related genes are located in close proximity to each other on chromosome 20p13. Multiple alternatively spliced transcript variants have been determined for this gene. [provided by RefSeq
Other Designations	OTTHUMP0000030001 SHP substrate-1 brain-immunoglobulin-like molecule with tyrosine-base d activation motifs macrophage fusion receptor myd-1 antigen protein tyrosine phosphatase, non-receptor type substrate 1 signal regulatory protein, alpha type 1 signal