


PTK2 polyclonal antibody

Catalog # PAB16301 Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human breast with PTK2 polyclonal antibody (Cat # PAB16301).

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic peptide of PTK2.
Immunogen	A synthetic peptide (conjugated with KLH) corresponding to human PTK2.
Host	Rabbit
Reactivity	Bovine, Chicken, Dog, Hamster, Human, Monkey, Mouse, Rat
Specificity	Kinase domain of human.
Form	Liquid
Purification	Immunoaffinity purification
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (13 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

😵 Abnova

Product Information

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human breast with PTK2 polyclonal antibody (Cat # PAB16301).

Gene Info — PTK2

Entrez GenelD	<u>5747</u>
Protein Accession#	<u>Q05397</u>
Gene Name	PTK2
Gene Alias	FADK, FAK, FAK1, pp125FAK
Gene Description	PTK2 protein tyrosine kinase 2
Omim ID	<u>600758</u>
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. Th e encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks signific ant sequence similarity to kinases from other subfamilies. Activation of this gene may be an impor tant early step in cell growth and intracellular signal transduction pathways triggered in response t o certain neural peptides or to cell interactions with the extracellular matrix. At least four transcript variants encoding four different isoforms have been found for this gene, but the full-length natures of only two of them have been determined. [provided by RefSeq

Pathway

- Axon guidance
- <u>Chemokine signaling pathway</u>
- ErbB signaling pathway

😵 Abnova

- Focal adhesion
- Leukocyte transendothelial migration
- Pathways in cancer
- <u>Regulation of actin cytoskeleton</u>
- Small cell lung cancer
- VEGF signaling pathway

Disease

- <u>Autistic Disorder</u>
- Genetic Predisposition to Disease
- HIV Infections
- Leukemia
- Mental Retardation
- <u>Neovascularization</u>
- <u>Psychotic Disorders</u>
- Schizophrenia