


GRM5 polyclonal antibody

Catalog # PAB16484

Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human colon (myenteric plexus) with GRM5 polyclonal antibody (Cat # PAB16484).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of GRM5.
Immunogen	A synthetic peptide (conjugated with KLH) corresponding to human GRM5.
Host	Rabbit
Reactivity	Bovine, Chicken, Dog, Hamster, Horse, Human, Monkey, Mouse, Pig, Rabbit, Rat
Specificity	N-terminal extracellular domain of human.
Form	Liquid
Purification	Immunoaffinity purification
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (5.6 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human colon (myenteric plexus) with GRM5 polyclonal antibody (Cat # PAB16484).

Gene Info — GRM5

Entrez GeneID [2915](#)

Protein Accession# [P41594](#)

Gene Name GRM5

Gene Alias GPRC1E, MGLUR5, mGlu5

Gene Description glutamate receptor, metabotropic 5

Omim ID [604102](#)

Gene Ontology [Hyperlink](#)

Gene Summary

L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations -

Pathway

- [Calcium signaling pathway](#)
- [Gap junction](#)

- [Long-term depression](#)
- [Long-term potentiation](#)
- [Neuroactive ligand-receptor interaction](#)

Disease

- [Alcoholism](#)
- [Anorexia Nervosa](#)
- [Bulimia](#)
- [Cardiovascular Diseases](#)
- [Cognition](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Mental Disorders](#)
- [Schizophrenia](#)
- [Schizophrenic Psychology](#)
- [Tobacco Use Disorder](#)
- [Weight Gain](#)