


FLT1 polyclonal antibody

Catalog # PAB16509

Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human placenta (fetal capillary endothelial cells) with FLT1 polyclonal antibody (Cat # PAB16509).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of FLT1.
Immunogen	A synthetic peptide (conjugated with KLH) corresponding to human FLT1.
Host	Rabbit
Reactivity	Human
Specificity	N-terminal domain of human.
Form	Liquid
Purification	Immunoaffinity purification
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (5-10 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human placenta (fetal capillary endothelial cells) with FLT1 polyclonal antibody (Cat # PAB16509).

Gene Info — FLT1

Entrez GeneID [2321](#)

Protein Accession# [P17948](#)

Gene Name FLT1

Gene Alias FLT, VEGFR1

Gene Description fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor)

Omim ID [165070](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia

Other Designations fms-related tyrosine kinase 1|soluble VEGF receptor 1-14|soluble VEGFR1 variant 2|soluble VEGFR1 variant 21|vascular endothelial growth factor/vascular permeability factor receptor

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)

- [Focal adhesion](#)

Disease

- [Abortion](#)
- [Adenocarcinoma](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Cardiovascular Diseases](#)
- [Cell Transformation](#)
- [Chorioamnionitis](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Growth Retardation](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Hypercholesterolemia](#)
- [Hypersensitivity](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lymphoma](#)
- [Malaria](#)
- [Melanoma](#)
- [Neovascularization](#)

- [Obstetric Labor](#)
- [Placenta Diseases](#)
- [Pre-Eclampsia](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Sarcoidosis](#)
- [Scleroderma](#)
- [Skin Neoplasms](#)
- [Vaginosis](#)