


TRIP13 polyclonal antibody

Catalog # PAB17390 Size 100 ug

Applications



Western Blot (Cell lysate)

Western blot analysis of extracts from LoVo cells (Lane 1) and NIH/3T3 cells (Lane 2 and lane 3), using TRIP13 polyclonal antibody (Cat # PAB17390). Peptide "+" means "with peptide blocking".

Specification

Product Description Rabbit polyclonal antibody raised against synthetic peptide of TRIP13.

Immunogen A synthetic peptide corresponding to C-terminus of human TRIP13.

Host Rabbit

Reactivity Human, Mouse, Rat

Specificity This antibody detects endogenous levels of total TRIP13 protein.

Form Liquid

Recommend Usage Western Blot (1:500-1:1000)
ELISA (1:5000)
The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, pH 7.4 (150mM NaCl, 0.02% sodium azide, 50% glycerol)

Storage Instruction Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of extracts from LoVo cells (Lane 1) and NIH/3T3 cells (Lane 2 and lane 3), using TRIP13 polyclonal antibody (Cat # PAB17390).

Peptide "+" means "with peptide blocking".

Gene Info — TRIP13

Entrez GeneID	9319
---------------	----------------------

Protein Accession#	Q15645
--------------------	------------------------

Gene Name	TRIP13
-----------	--------

Gene Alias	16E1BP
------------	--------

Gene Description	thyroid hormone receptor interactor 13
------------------	--

Omim ID	604507
---------	------------------------

Gene Ontology	Hyperlink
---------------	---------------------------

Other Designations	thyroid receptor interacting protein 13
--------------------	---

Publication Reference

- [Immunoaffinity profiling of tyrosine phosphorylation in cancer cells.](#)

Rush J, Moritz A, Lee KA, Guo A, Goss VL, Spek EJ, Zhang H, Zha XM, Polakiewicz RD, Comb MJ.

Nature Biotechnology 2005 Jan; 23(1):94.

- [The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection \(MGC\).](#)

Gerhard DS, Wagner L, Feingold EA, Shenmen CM, Grouse LH, Schuler G, Klein SL, Old S, Rasooly R, Good P, Guyer M, Peck AM, Derge JG, Lipman D, Collins FS, Jang W, Sherry S, Feolo M, Misquitta L, Lee E, Rotmistrovsky K, Greenhut SF, Schaefer CF, Buetow K, Bonner TI, Haussler D, Kent J, Kiekhaus M, Furey T, Brent M, Prange C, Schreiber K, Shapiro N, Bhat NK, Hopkins RF, Hsie F, Driscoll T, Soares MB, Casavant TL, Scheetz TE, Brownstein MJ, Usdin TB, Toshiyuki S, Carninci P, Piao Y, Dudekula DB, K

Genome Research 2004 Oct; 14(10B):2121.

- [Two classes of human papillomavirus type 16 E1 mutants suggest pleiotropic conformational constraints affecting E1 multimerization, E2 interaction, and interaction with cellular proteins.](#)

Yasugi T, Vidal M, Sakai H, Howley PM, Benson JD.

Journal of Virology 1997 Aug; 71(8):5942.