


SQSTM1 polyclonal antibody

Catalog # PAB1750


Size 400 uL

Applications




Western Blot (Transfected lysate)

Western blot analysis of SQSTM1 (arrow) using SQSTM1 polyclonal antibody (Cat # PAB1750). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the SQSTM1 gene (Lane 2) (Origene Technologies).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human hepatocellular carcinoma tissue reacted with SQSTM1 polyclonal antibody (Cat # PAB1750), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. HC = hepatocarcinoma. Data courtesy of Dr. Eeva-Liisa Eskelinen, University of Helsinki, Finland.



Immunofluorescence

Immunofluorescence staining of SQSTM1 polyclonal antibody (Cat # PAB1750) on Methanol-fixed and PFA fixed HeLa cells. Data courtesy of Dr. Eeva-Liisa Eskelinen, University of Helsinki, Finland.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of SQSTM1.
Immunogen	A synthetic peptide (conjugated with KLH) corresponding to C-terminus of human SQSTM1.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Protein G purification
Recommend Usage	Western Blot (1:1000) Immunohistochemistry (1:50-100) Immunofluorescence (1:50-100) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Transfected lysate)

Western blot analysis of SQSTM1 (arrow) using SQSTM1 polyclonal antibody (Cat # PAB1750). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the SQSTM1 gene (Lane 2) (Origene Technologies).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human hepatocellular carcinoma tissue reacted with SQSTM1 polyclonal antibody (Cat # PAB1750), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. HC = hepatocarcinoma. Data courtesy of Dr. Eeva-Liisa Eskelinen, University of Helsinki, Finland.

- Immunofluorescence

Immunofluorescence staining of SQSTM1 polyclonal antibody (Cat # PAB1750) on Methanol-fixed and PFA fixed HeLa cells. Data courtesy of Dr. Eeva-Liisa Eskelinen, University of Helsinki, Finland.

Gene Info — SQSTM1

Entrez GeneID	8878
Protein Accession#	NP_003891;Q13501
Gene Name	SQSTM1
Gene Alias	A170, OSIL, PDB3, ZIP3, p60, p62, p62B
Gene Description	sequestosome 1
Omim ID	601530 602080
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF-κB) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-κB in response to upstream signals. Alternatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone. [provided by RefSeq]
Other Designations	EBI3-associated protein p60 Paget disease of bone 3 oxidative stress induced like phosphotyrosine independent ligand for the Lck SH2 domain p62 ubiquitin-binding protein p62

Publication Reference

- [Inhibition of Autophagic Turnover in β-cells by Fatty acids and Glucose leads to Apoptotic cell Death.](#)
 Mir SU, George NM, Zahoor L, Harms R, Guinn Z, Sarvetnick NE.
 The Journal of Biological Chemistry 2015 Mar; 290(10):6071.
 Application: IF, WB-Ce, Human, Mouse, MIN6, INS-1 cells, Pancreatic islets
- [Absence of autophagy results in reactive oxygen species-dependent amplification of RLR signaling.](#)
 Tal MC, Sasai M, Lee HK, Yordy B, Shadel GS, Iwasaki A.
 PNAS 2009 Feb; 106(8):2770.
- [Impaired protein aggregate handling and clearance underlie the pathogenesis of p97/VCP-associated disease.](#)
 Ju JS, Miller SE, Hanson PI, Weihl CC.
 The Journal of Biological Chemistry 2008 Aug; 283(44):30289.

Disease

- [Genetic Predisposition to Disease](#)

- [Multiple System Atrophy](#)
- [Osteitis Deformans](#)
- [Tobacco Use Disorder](#)